

# A short proof of Mader's $\mathcal{S}$ -paths theorem

Alexander Schrijver<sup>1</sup>

**Abstract.** For an undirected graph  $G = (V, E)$  and a collection  $\mathcal{S}$  of disjoint subsets of  $V$ , an  $\mathcal{S}$ -path is a path connecting different sets in  $\mathcal{S}$ . We give a short proof of Mader's min-max theorem for the maximum number of disjoint  $\mathcal{S}$ -paths.

Let  $G = (V, E)$  be an undirected graph and let  $\mathcal{S}$  be a collection of disjoint subsets of  $V$ . An  $\mathcal{S}$ -path is a path connecting two different sets in  $\mathcal{S}$ . Mader [4] gave the following min-max relation for the maximum number of (vertex-)disjoint  $\mathcal{S}$ -paths, where  $S := \bigcup \mathcal{S}$ .

**Mader's  $\mathcal{S}$ -paths theorem.** *The maximum number of disjoint  $\mathcal{S}$ -paths is equal to the minimum value of*

$$(1) \quad |U_0| + \sum_{i=1}^n \lfloor \frac{1}{2} |B_i| \rfloor,$$

taken over all partitions  $U_0, \dots, U_n$  of  $V$  such that each  $\mathcal{S}$ -path disjoint from  $U_0$ , traverses some edge spanned by some  $U_i$ . Here  $B_i$  denotes the set of vertices in  $U_i$  that belong to  $S$  or have a neighbour in  $V \setminus (U_0 \cup U_i)$ .

Lovász [3] gave an alternative proof, by deriving it from his matroid matching theorem. Here we give a short proof of Mader's theorem.

Let  $\mu$  be the minimum value obtained in (1). Trivially, the maximum number of disjoint  $\mathcal{S}$ -paths is at most  $\mu$ , since any  $\mathcal{S}$ -path disjoint from  $U_0$  and traversing an edge spanned by  $U_i$ , traverses at least two vertices in  $B_i$ .

I. First, the case where  $|T| = 1$  for each  $T \in \mathcal{S}$  was shown by Gallai [2], by reduction to matching theory as follows. Let the graph  $\tilde{G} = (\tilde{V}, \tilde{E})$  arise from  $G$  by adding a disjoint copy  $G'$  of  $G - S$ , and making the copy  $v'$  of each  $v \in V \setminus S$  adjacent to  $v$  and to all neighbours of  $v$  in  $G$ . We claim that  $\tilde{G}$  has a matching of size  $\mu + |V \setminus S|$ . Indeed, by the Tutte-Berge formula ([5],[1]), it suffices to prove that for any  $\tilde{U}_0 \subseteq \tilde{V}$ :

$$(2) \quad |\tilde{U}_0| + \sum_{i=1}^n \lfloor \frac{1}{2} |\tilde{U}_i| \rfloor \geq \mu + |V \setminus S|,$$

where  $\tilde{U}_1, \dots, \tilde{U}_n$  are the components of  $\tilde{G} - \tilde{U}_0$ . Now if for some  $v \in V \setminus S$  exactly one of  $v, v'$  belongs to  $\tilde{U}_0$ , then we can delete it from  $\tilde{U}_0$ , thereby not increasing the left hand side of (2). So we can assume that for each  $v \in V \setminus S$ , either  $v, v' \in \tilde{U}_0$  or  $v, v' \notin \tilde{U}_0$ . Let  $U_i := \tilde{U}_i \cap V$  for  $i = 0, \dots, n$ . Then  $U_1, \dots, U_n$  are the components of  $G - U_0$ , and we have:

$$(3) \quad |\tilde{U}_0| + \sum_{i=1}^n \lfloor \frac{1}{2} |\tilde{U}_i| \rfloor = |U_0| + \sum_{i=1}^n \lfloor \frac{1}{2} |U_i \cap S| \rfloor + |V \setminus S| \geq \mu + |V \setminus S|$$

(since in this case  $B_i = U_i \cap S$  for  $i = 1, \dots, n$ ), showing (2).

---

<sup>1</sup>CWI, Kruislaan 413, 1098 SJ Amsterdam, The Netherlands, and Department of Mathematics, University of Amsterdam, Plantage Muidergracht 24, 1018 TV Amsterdam, The Netherlands.

So  $\tilde{G}$  has a matching  $M$  of size  $\mu + |V \setminus S|$ . Let  $N$  be the matching  $\{vv' | v \in V \setminus S\}$  in  $\tilde{G}$ . As  $|M| = \mu + |V \setminus S| = \mu + |N|$ , the union  $M \cup N$  has at least  $\mu$  components with more edges in  $M$  than in  $N$ . Each such component is a path connecting two vertices in  $S$ . Then contracting the edges in  $N$  yields  $\mu$  disjoint  $\mathcal{S}$ -paths in  $G$ .

II. We now consider the general case. Fixing  $V$ , choose a counterexample  $E, \mathcal{S}$  minimizing

$$(4) \quad |E| - |\{\{t, u\} | t, u \in V, \exists T, U \in \mathcal{S} : t \in T, u \in U, T \neq U\}|.$$

By part I, there exists a  $T \in \mathcal{S}$  with  $|T| \geq 2$ . Then  $T$  is independent in  $G$ , since any edge  $e$  spanned by  $T$  can be deleted without changing the maximum and minimum value in Mader's theorem (as any  $\mathcal{S}$ -path traversing  $e$  contains an  $\mathcal{S}$ -path not containing  $e$ , and as deleting  $e$  does not change any set  $B_i$ ), while decreasing (4).

Choose  $s \in T$ . Replacing  $\mathcal{S}$  by  $\mathcal{S}' := (\mathcal{S} \setminus \{T\}) \cup \{T \setminus \{s\}, \{s\}\}$  decreases (4), but not the minimum in Mader's theorem (as each  $\mathcal{S}$ -path is an  $\mathcal{S}'$ -path and as  $\bigcup \mathcal{S}' = S$ ). So there exists a collection  $\mathcal{P}$  of  $\mu$  disjoint  $\mathcal{S}'$ -paths. We can assume that no path in  $\mathcal{P}$  has any internal vertex in  $S$ .

Necessarily, there is a path  $P_0 \in \mathcal{P}$  connecting  $s$  with another vertex in  $T$ , all other paths in  $\mathcal{P}$  being  $\mathcal{S}$ -paths. Let  $u$  be an internal vertex of  $P_0$ . Replacing  $\mathcal{S}$  by  $\mathcal{S}'' := (\mathcal{S} \setminus \{T\}) \cup \{T \cup \{u\}\}$  decreases (4), but not the minimum in Mader's theorem (as each  $\mathcal{S}$ -path is an  $\mathcal{S}''$ -path and as  $\bigcup \mathcal{S}'' \supset S$ ). So there exists a collection  $\mathcal{Q}$  of  $\mu$  disjoint  $\mathcal{S}''$ -paths. Choose  $\mathcal{Q}$  such that no internal vertex of any path in  $\mathcal{Q}$  belongs to  $S \cup \{u\}$ , and such that  $\mathcal{Q}$  uses a minimal number of edges not used by  $\mathcal{P}$ .

Necessarily,  $u$  is an end of some path  $Q_0 \in \mathcal{Q}$ , all other paths in  $\mathcal{Q}$  being  $\mathcal{S}$ -paths. As  $|\mathcal{P}| = |\mathcal{Q}|$  and as  $u$  is not an end of any path in  $\mathcal{P}$ , there exists an end  $v$  of some path  $P \in \mathcal{P}$  that is not an end of any path in  $\mathcal{Q}$ . Now  $P$  intersects at least one path in  $\mathcal{Q}$  (since otherwise  $P \neq P_0$ , and  $(\mathcal{Q} \setminus \{Q_0\}) \cup \{P\}$  would consist of  $\mu$  disjoint  $\mathcal{S}$ -paths). So when following  $P$  starting at  $v$ , there is a first vertex  $w$  that is on some path in  $\mathcal{Q}$ , say on  $Q \in \mathcal{Q}$ .

For any end  $x$  of  $Q$  let  $Q^x$  be the  $x - w$  part of  $Q$ , let  $P^v$  be the  $v - w$  part of  $P$ , and let  $U$  be the set in  $\mathcal{S}''$  containing  $v$ . Then for any end  $x$  of  $Q$  we have that  $Q^x$  is part of  $P$  or the other end of  $Q$  belongs to  $U$ , since otherwise by rerouting part  $Q^x$  of  $Q$  along  $P^v$ ,  $Q$  remains an  $\mathcal{S}''$ -path disjoint from the other paths in  $\mathcal{Q}$ , while we decrease the number of edges used by  $\mathcal{Q}$  and not by  $\mathcal{P}$ , contradicting the minimality assumption.

Let  $y, z$  be the ends of  $Q$ . We can assume that  $y \notin U$ . Then  $Q^y$  is part of  $P$ , hence  $Q^y$  is not part of  $P$  (as  $Q$  is not part of  $P$ , as otherwise  $Q = P$ , and hence  $v$  is an end of  $Q$ ), so  $z \in U$ . As  $z$  is on  $P$  and as also  $v$  belongs to  $U$  and is on  $P$ , we have  $P = P_0$ . So  $U = T \cup \{u\}$  and  $Q = Q_0$  (since  $Q^z$  is part of  $P$ , so  $z = u$ ). But then rerouting part  $Q^z$  of  $Q$  along  $P^v$  gives  $\mu$  disjoint  $\mathcal{S}$ -paths, contradicting our assumption.

## References

- [1] C. Berge, Sur le couplage maximum d'un graphe, *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences [Paris]* 247 (1958) 258–259.
- [2] T. Gallai, Maximum-minimum Sätze und verallgemeinerte Faktoren von Graphen, *Acta Mathematica Academiae Scientiarum Hungaricae* 12 (1961) 131–173.
- [3] L. Lovász, Matroid matching and some applications, *Journal of Combinatorial Theory, Series B* 28 (1980) 208–236.

- [4] W. Mader, Über die Maximalzahl kreuzungsfreier  $H$ -Wege, *Archiv der Mathematik (Basel)* 31 (1978) 387–402.
- [5] W.T. Tutte, The factorization of linear graphs, *The Journal of the London Mathematical Society* 22 (1947) 107–111.