

RESEARCH ARTICLE

Laplacian Instability of Planar Streamer Ionization Fronts—An Example of Pulled Front Analysis

Gianne Derks · Ute Ebert · Bernard Meulenbroek

Received: 15 June 2007 / Accepted: 21 March 2008 / Published online: 4 June 2008
© Springer Science+Business Media, LLC 2008

Abstract Streamer ionization fronts are pulled fronts that propagate into a linearly unstable state; the spatial decay of the initial condition of a planar front selects dynamically one specific long-time attractor out of a continuous family. A stability analysis for perturbations in the transverse direction has to take these features into account. In this paper we show how to apply the Evans function in a weighted space

Communicated by G. Gottwald.

G. Derks acknowledges a travel grant of the Royal Society, which initiated this research, and a visitor grant of the Dutch funding agency NWO and the NWO-mathematics cluster NDNS⁺ to finish the work. The work was also supported by a CWI PhD grant for B. Meulenbroek.

G. Derks (✉)

Department of Mathematics, University of Surrey, Guildford, Surrey, GU2 7XH, UK
e-mail: G.Derks@surrey.ac.uk

U. Ebert · B. Meulenbroek

Center for Mathematics and Computer Science (CWI), P.O. Box 94079, 1090 GB Amsterdam, The Netherlands

U. Ebert

e-mail: ebert@cwi.nl

U. Ebert

Department of Applied Physics, Eindhoven University of Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

B. Meulenbroek

Faculty of Electrical Engineering, Mathematics and Computer Science, Delft Univ. Techn., P.O. Box 5031, 2600 GA Delft, The Netherlands
e-mail: B.J.Meulenbroek@tudelft.nl

for this stability analysis. Zeros of the Evans function indicate the intersection of the stable and unstable manifolds; they are used to determine the eigenvalues. Within this Evans function framework, we define a numerical dynamical systems method for the calculation of the dispersion relation as an eigenvalue problem. We also derive dispersion curves for different values of the electron diffusion constant and of the electric field ahead of the front. Numerical solutions of the initial value problem confirm the eigenvalue calculations. The numerical work is complemented with an analysis of the Evans function leading to analytical expressions for the dispersion relation in the limit of small and large wave numbers. The paper concludes with a fit formula for intermediate wave numbers. This empirical fit supports the conjecture that the smallest unstable wave length of the Laplacian instability is proportional to the diffusion length that characterizes the leading edge of the pulled ionization front.

Keywords Pulled front · Stability analysis · Streamer ionization front · Dispersion relation · Wave selection of Laplacian instability

Mathematics Subject Classification (2000) 37L15 · 34L16 · 35Q99

1 Introduction

1.1 The Streamer Phenomenon, Ionization Fronts and Laplacian Instability

A streamer is the first stage of electric breakdown in large volumes. It paves the way for sparks and lightning, but also occurs without successive breakdown in phenomena like sprite discharges above thunderclouds or in corona discharges used in numerous technical applications. Recent reviews of relevant phenomena can be found in Ebert et al. (2006) and Starikovskaia (2006). Considered as a nonlinear phenomenon, the streamer is a finger-shaped ionized region that propagates by self-generated field enhancement at its tip into nonionized media. It has multiple scales as described in Ebert et al. (2006); as a consequence one can investigate a hierarchy of models on different levels of refinement that are reductions of each other. One can start from the reduction from a particle to a continuum model (Li et al. 2007) to the reduction from a continuum model to a moving boundary model (Brau et al. 2008) up to the formulation of effective models for complete multiple-branched streamer trees without inner structure. These trees are known as “dielectric breakdown models” (Niemeyer et al. 1984, 1989; Pasko et al. 2001; Bawagan 1997). All these reductions are the subject of current research; the present paper analyzes the stability of fronts in the continuum model; the resulting dispersion relation provides a test case for moving boundary approximations.

Specifically, simulations of the simplest continuum model for negative streamers (Dhali and Williams 1985, 1987; Vitello et al. 1994) have established the formation of a thin boundary layer around the streamer head. This layer is an ionization front that also carries a net negative electric charge. (Positive streamers with positive net charge occur as well, but are not the subject of the present study.) The configuration of the charge in a thin layer leads to the above-mentioned field enhancement at the streamer head that creates high ionization rates and electron drift velocities and hence lets the streamer rapidly penetrate the nonionized region. More recent numerical in-

vestigations show that the boundary layer or front can undergo a Laplacian instability that lets the streamer branch (Arrayás et al. 2002; Rocco et al. 2002; Montijn et al. 2006a, 2006b). (We remark that an additional interaction mechanism in composite gases like air somewhat modifies this picture (Luque et al. 2007) while the present analysis applies to negative streamers in simple gases like pure nitrogen or argon.)

1.2 Moving Boundary Layers and the Transversal Instability of Pulled Fronts

The streamer can be considered as a phenomenon where an ionized phase is separated from a nonionized phase by a moving thin front. This concept (Ebert et al. 1996; Arrayás et al. 2002) implies that streamers show similar features as moving boundary problems like viscous fingers, solidification fronts propagating into undercooled liquids, growth of bacterial colonies or corals in a diffusive field of food etc. Quantitative predictions within such models require a proper understanding of the front dynamics, in particular, of their stability against perturbations in the transversal direction. This stability determines whether perturbations of the front position will grow or shrink, and in the long term whether the streamer will branch or not. As a first insight, one would therefore like to analyze the stability of planar fronts against transversal perturbations, more specifically, the growth or shrinking rate $s(k)$ of a linear perturbation with transversal wavelength $2\pi/k$.

The ionization front in the model for a negative streamer in a pure gas as treated in Dhali and Williams (1985, 1987), Vitello et al. (1994), Ebert et al. (1996, 1997), Arrayás et al. (2002), Rocco et al. (2002), Montijn et al. (2006a, 2006b), Brau et al. (2008), including electron diffusion, creates a so-called pulled front that has a number of peculiar mathematical properties: (i) for each velocity $v \geq v^*$, there is a dynamically stable front solution where the stability is conditional on the spatial decay of the perturbation, hence the long time dynamics is selected by the spatial decay of the initial front for $z \rightarrow \infty$ (where z is the spatial variable along the front); (ii) the convergence toward this front is algebraically slow in time (Ebert and van Saarloos 1998, 2000a); (iii) these slow dynamics are determined in the leading edge of the front that, in principle, extends up to $z \rightarrow \infty$ and in the dynamically relevant space it will cause Fredholm integrals in the linear stability analysis to diverge, therefore curvature corrections cannot be calculated in the established manner (Ebert and van Saarloos 2000b); (iv) the unconventional location of the dynamically relevant region ahead of the front also requires particular care in numerical solutions with adaptive grid refinement (Montijn et al. 2006b). For the calculation of the dispersion relation, which can be phrased as an eigenvalue problem for $s(k)$, these features pose two challenges: first, the condition on the one-dimensional dynamical stability and algebraic convergence properties, which are typical for pulled fronts, will lead to an apparently degenerate eigenvalue problem. Second, in a neighborhood of the origin, the dispersion curve $s(k)$ is near the continuous spectrum. Hence numerical calculations of the eigenvalue problem with finite difference, collocation or spectral methods often lead to spurious eigenvalues. A dynamical systems method involving stable and unstable manifolds avoids this problem. The stable and unstable manifolds are at least two-dimensional and an exterior algebra approach is employed to calculate the manifolds accurately.

In Ebert and Arrayás (2001), Arrayás et al. (2002), Arrayás and Ebert (2004), the treatment of pulled fronts and more-dimensional stable/unstable manifolds was circumvented by neglecting the electron diffusion that acts as a singular perturbation. In this way, the leading edge of the front together with its mathematical challenges is removed and the eigenvalue problem can be solved using shooting on the one-dimensional stable/unstable manifolds. The resulting problem is characterized by two length scales, namely the length scale $2\pi/k$ of the transversal perturbation, and the longitudinal length scale of electric screening through the front that will be denoted by ℓ_α . The dispersion relation in this case shows a quite unconventional behavior, namely a short wavelength instability whose consequences are further investigated in Meulenbroek et al. (2005) and Ebert et al. (2007). In the present paper, we analyze the dispersion relation including diffusion, mastering the above challenges and deriving quantitative results through a combination of analytical and numerical methods.

1.3 The Evans Function and Pulled Fronts

The Evans function is an analytic function whose zeros correspond to the eigenvalues of a spectral problem, usually a linearization about a coherent structure like a front or solitary wave. It was first introduced in Evans (1975) and generalized in Alexander et al. (1990). In the last decade, the Evans function has been applied in the context of many problems and various extensions and generalizations have been found, see the review papers (Kapitula and Sandstede 2004; Sandstede 2002) and references therein. One of the first uses of the Evans function in the analysis of a planar front can be found in Terman (1990), which analyzes the stability of a planar wave in a reaction diffusion system arising in a combustion model. In the current paper we show how pulled fronts can be analyzed with the Evans function by using a definition of the Evans function in a weighted space.

To define the Evans function, one writes the eigenvalue problem as a linear, first-order, dynamical system with respect to the spatial variable z . Along the dispersion curve $s(k)$, the dynamical system has a solution that is bounded for all values of z . This can be phrased in a more dynamical way as: the manifold of solutions that are exponentially decaying for $z \rightarrow +\infty$ (stable manifold) and the manifold of solutions that are exponentially decaying for $z \rightarrow -\infty$ (unstable manifold) have a nontrivial intersection along the dispersion curve. The Evans function is a function of the spectral parameters s and k , which vanishes if the stable and unstable manifolds have a nontrivial intersection. Hence the Evans function can be viewed as a Melnikov function or a Wronskian determinant, see also Kapitula (1999) or references in there.

In case of a pulled front, the definition of the stable manifold, and hence the Evans function, is not straightforward. The temporal stability of the asymptotic state of the pulled front at $+\infty$ is conditional on the spatial decay of the perturbation. So this decay condition should be included in the definition of the stable manifold, otherwise the dimension of this manifold might be too large. We will show that this condition can be built in the definition of the stable manifold by considering the stable manifold in a weighted space. The Evans function is defined by using the weighted space for the stable manifold. Hence the dispersion curve $s(k)$ can be found as a curve of zeros of this Evans function.

1.4 Organization of the Paper

In Sect. 2, we recall the model equations and the construction and properties of planar fronts. In particular, we summarize the multiplicity, stability, dynamical selection and convergence rate of these pulled fronts. In Sect. 3, the stability of these fronts is investigated as an eigenvalue problem for the dispersion relation $s(k)$ of a linear perturbation with wave number k . The dispersion relation depends on the far electric field E_∞ and the electron diffusion D as external parameters. In the stability analysis of the pulled ionization fronts, a constraint is imposed on the asymptotic spatial decay rate of the perturbations. This constraint corresponds to the decay condition for the one-dimensional stability, but has to be chosen quite subtly to avoid problems with the algebraic decay of the front solution. A consequence of the decay condition is that the eigenvalue problem (dispersion relation) is solved in a weighted space. In this weighted space, the apparent degeneracies have disappeared, the stable and unstable manifolds of the ODE related to the eigenvalue problem are well defined and intersections of those manifolds are determined by using the Evans function. In Sect. 3.4, dispersion relations for positive s are derived numerically for a number of pairs of external parameters (E_∞, D) . The numerical implementation of the Evans function uses exterior algebra to reliably solve for the higher dimensional stable and unstable manifolds. In Sect. 4, the numerical dispersion relation is tested thoroughly and confirmed with numerical simulations of the initial value problem for the complete PDE model for the particular values $(E_\infty, D) = (-1, 0.1)$ where $D = 0.1$ is typically used for nitrogen (Dhali and Williams 1985, 1987; Vitello et al. 1994; Ebert et al. 1996, 1997; Arrayás et al. 2002; Rocco et al. 2002; Montijn et al. 2006a, 2006b) and $E_\infty = -1$ is a representative value for the electric field. The later sections treat either general (E_∞, D) analytically or a larger range of (E_∞, D) numerically.

In Sect. 5, explicit analytical asymptotic relations for the dispersion relation $s(k)$ are derived for the limits of small and large wave numbers k . For $k = 0$, two explicit eigenfunctions are known (which are related to the translation and gauge symmetry in the problem). These explicit solutions lead to expressions for the solutions on the stable manifold for small wave numbers. The interaction between the slow and fast behavior on this manifold leads to an asymptotic dispersion relation for small k . For large wave numbers, the eigenvalue problem for the dispersion relation is dominated by a constant coefficient eigenvalue problem. An eigenvalue exists only if this constant coefficient system has no spectral gap. Using exponential dichotomies and the roughness theorem, the asymptotics of the dispersion relation is derived by a contradiction argument. In Sect. 6, these asymptotic limits are tested on the numerical data derived in Sect. 3. It is found that the asymptotic limit for small k fits the data very well, while the asymptotic limit for large k is not yet applicable in the range where $s(k)$ is positive. After a discussion of relevant physical scales, we suggest a fit formula joining the analytical small k asymptotic limit with our physically motivated guess. This formula fits the numerical data well for practical purposes and strongly supports the conjecture that the smallest unstable wavelength is proportional to the diffusion length that determines the leading edge of the pulled front.

2 The Streamer Model and Its Ionization Fronts

In this section we describe the streamer model and summarize the features of planar ionization fronts as solutions of the purely one-dimensional model as a preparation for the stability analysis in the dimensions transversal to the front. In particular, we recall the multiplicity of the front solutions that penetrate a linearly dynamically unstable state, and the dynamical selection of the pulled front.

2.1 Model Equations

We investigate negative fronts within the minimal streamer model, i.e., within a “fluid approximation” with local field-dependent impact ionization reaction in a nonattaching gas like argon or nitrogen (Ebert et al. 1996, 1997; Ebert and Arrayás 2001; Arrayás et al. 2002; Rocco et al. 2002). The equations for this model in dimensionless quantities are

$$\partial_t \sigma - D \nabla^2 \sigma - \nabla \cdot (\sigma \mathbf{E}) = \sigma f(|\mathbf{E}|), \quad (2.1)$$

$$\partial_t \rho = \sigma f(|\mathbf{E}|), \quad (2.2)$$

$$\nabla \cdot \mathbf{E} = \rho - \sigma, \quad \mathbf{E} = -\nabla \phi, \quad (2.3)$$

where σ is the electron and ρ the ion density, \mathbf{E} is the electric field and ϕ is the electrostatic potential. For physical parameters and dimensional analysis, we refer to discussions in Ebert et al. (1996, 1997), Ebert and Arrayás (2001), Arrayás et al. (2002), Rocco et al. (2002). The electron current is approximated by diffusion and advection $-D \nabla \sigma - \sigma \mathbf{E}$. The ion current is neglected because the front dynamics takes place on the fast time scale of the electrons and the ion mobility is much smaller. Electron–ion pairs are assumed to be generated with rate $\sigma f(|\mathbf{E}|) = \sigma |\mathbf{E}| \alpha(|\mathbf{E}|)$, where $\sigma |\mathbf{E}|$ is the absolute value of electron drift current and $\alpha(|\mathbf{E}|)$ the effective impact ionization cross-section within a field \mathbf{E} . Hence $f(|\mathbf{E}|)$ is

$$f(|\mathbf{E}|) = |\mathbf{E}| \alpha(|\mathbf{E}|). \quad (2.4)$$

For numerical calculations, we use the Townsend approximation $\alpha(|\mathbf{E}|) = e^{-1/|\mathbf{E}|}$ (Ebert et al. 1996, 1997; Ebert and Arrayás 2001; Arrayás et al. 2002; Rocco et al. 2002). For analytical calculations, an arbitrary function $\alpha(|\mathbf{E}|) \geq 0$ can be chosen where we assume that $\alpha(0) = 0$ and therefore $f(0) = 0 = f'(0)$. We will furthermore assume that $\alpha(|\mathbf{E}|)$ is monotonically increasing in $|\mathbf{E}|$, this is a sufficient criterion for the front to be a pulled one (Ebert and van Saarloos 2000a). The electric field can be calculated in electrostatic approximation $\mathbf{E} = -\nabla \phi$.

Mathematically, the model (2.1)–(2.3) describes the dynamics of the three scalar fields σ , ρ and ϕ . It is a set of reaction-advection-diffusion equations for the charged species σ and ρ coupled nonlinearly to the Poisson equation of electrostatics.

2.2 Two Types of Stationary States

It follows immediately from (2.1)–(2.3) that there can be two types of stationary states of the system, one characterized by $\sigma \equiv 0$ and the other by $\mathbf{E} \equiv 0$ (as $f(|\mathbf{E}|) = 0$ implies $|\mathbf{E}| = 0$).

The stationary state with $\sigma \equiv 0$ is the nonionized state. As the dynamics is only carried by the electrons σ , there is no temporal evolution for $\sigma \equiv 0$ even if the ion density ρ has an arbitrary spatial distribution. The electric field $\mathbf{E} = -\nabla\phi$ then is determined by the solution of the Poisson equation $-\nabla^2\phi = \rho$ and by the boundary conditions on ϕ . In certain ionization fronts in semiconductor devices (Rodin et al. 2002), it is essential that the equivalent of ρ does not vanish in the nonionized region. In the gas discharges considered here, on the other hand, it is reasonable to assume that the nonionized initial state with $\sigma \equiv 0$ also has a vanishing ion density $\rho \equiv 0$, and therefore no space charges.

The stationary state with vanishing electric field $\mathbf{E} \equiv 0$ describes the ionized, electrically screened charge neutral plasma region behind an ionization front, the interior of the streamer. From $\mathbf{E} \equiv 0$ the identity $\nabla \cdot \mathbf{E} = 0$ follows immediately, and therefore electron and ion densities have to be equal $\sigma = \rho$. In the absence of a field, the electrons diffuse $\partial_t\sigma = D\nabla^2\sigma$ while the ions stay put $\partial_t\rho = 0$. Therefore, these densities can stay equal only if $\nabla^2\rho = 0$. Simulations (Dhali and Williams 1985; Dhali and Williams 1987; Vitello et al. 1994; Ebert et al. 1996; Ebert et al. 1997; Arrayás et al. 2002; Rocco et al. 2002; Montijn et al. 2006a; Montijn et al. 2006b) show that this occurs typically only if ρ is homogeneous (though counter examples can be constructed).

2.3 Planar Ionization Front Solutions

An ionization front separates such different outer regions: an electron-free and non-conducting state with an arbitrary electric field E_∞ ahead of the front from an ionized and electrically screened state with arbitrary, but equal density $\sigma^- = \rho^-$ of electrons and ions. In particular, we are interested in almost planar fronts propagating into a particle-free region $\rho = \sigma = 0$ (where therefore $\nabla^2\phi = 0$), and we study negative fronts, i.e., fronts with an electron surplus that propagate into the electron drift direction towards an asymptotic electric field $E_\infty < 0$. For a planar front, it follows from $\nabla^2\phi = -\nabla \cdot \mathbf{E} = 0$ that the electric field ahead of the front is homogeneous.

We assume that the front propagates into the positive z direction; the electric field ahead of a negative front then is $\mathbf{E} \rightarrow E_\infty \hat{\mathbf{z}}$, $E_\infty < 0$, for $z \rightarrow \infty$. (Here $\hat{\mathbf{z}}$ is the unit vector in the z -direction.) It is convenient to introduce the coordinate system $(x, y, \xi = z - vt)$ moving with the front velocity $\mathbf{v} = v\hat{\mathbf{z}}$. A planar, uniformly translating front is a stationary solution in this comoving frame, hence it depends only on the comoving coordinate ξ , and will be denoted by a lower index 0. A front satisfies

$$\begin{aligned} D\partial_\xi^2\sigma_0 + (v - \partial_\xi\phi_0)\partial_\xi\sigma_0 + \sigma_0(\rho_0 - \sigma_0) + \sigma_0 f_0 &= 0, \\ v\partial_\xi\rho_0 + \sigma_0 f_0 &= 0, \\ \partial_\xi^2\phi_0 + \rho_0 - \sigma_0 &= 0, \end{aligned} \quad (2.5)$$

where $f_0 = f(|E_0|)$. This system can be reduced to three first-order ordinary differential equations. First, due to electric gauge invariance, the system does not depend on ϕ_0 explicitly, but only on $E_0 = -\partial_\xi\phi_0$. Using the variable E_0 instead of ϕ_0 reduces the number of derivatives by one. Second, electric charge conservation

$\partial_t q + \nabla \cdot \mathbf{j} = 0$ can be rewritten in comoving coordinates for a uniformly translating front as $-v\partial_\xi q_0 + \partial_\xi j_0 = 0$. Therefore it can be integrated once $-vq_0 + j_0 = c$, $\partial_\xi c = 0$. In the present problem, the space charge is $q_0 = \rho_0 - \sigma_0$ and the electric current is $j_0 = -D\partial_\xi \sigma_0 - \sigma_0 E_0$. Furthermore, as there is a region with vanishing densities $\sigma_0 = 0 = \rho_0$ ahead of the front, the integration constant c vanishes in this region, and therefore everywhere. Thus the planar front (2.5) can be written as

$$\begin{aligned} D\partial_\xi \sigma_0 &= v(\rho_0 - \sigma_0) - E_0\sigma_0, \\ v\partial_\xi \rho_0 &= -\sigma_0 f(|E_0|), \\ \partial_\xi E_0 &= \rho_0 - \sigma_0, \end{aligned} \quad (2.6)$$

where $\partial_\xi \phi_0 = -E_0$ decouples from the other equations. The planar front equations imply that $E_0(\xi) < 0$ for all ξ when $E_\infty < 0$ (Ebert et al. 1997).

The fronts connect the states

$$\begin{pmatrix} \sigma_0 \\ \rho_0 \\ E_0 \end{pmatrix} \xrightarrow{\xi \rightarrow +\infty} \begin{pmatrix} 0 \\ 0 \\ E_\infty \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \sigma_0 \\ \rho_0 \\ E_0 \end{pmatrix} \xrightarrow{\xi \rightarrow -\infty} \begin{pmatrix} \sigma^- \\ \sigma^- \\ 0 \end{pmatrix}, \quad (2.7)$$

and the electrostatic potential ϕ_0 connects ϕ^- (for $\xi \rightarrow -\infty$) with $-E_\infty \xi + \phi^+$ (for $\xi \rightarrow +\infty$). The ionization density σ^- behind the front and the electrostatic potential difference $\phi^+ - \phi^-$ have to be determined for an arbitrarily chosen electric field E_∞ ahead of the front and for an arbitrary, but sufficiently large, front velocity v . (We remark that only the potential difference $\phi^+ - \phi^-$ matters due to the gauge invariance of the electrostatic potential as one easily verifies with the equations.) The fronts can be constructed as heteroclinic orbits in a three-dimensional space as demonstrated in Ebert et al. (1997).

The diffusion constant D is obviously a singular perturbation. For $D = 0$, the front equations can be solved analytically (Ebert et al. 1997; Arrayás and Ebert 2004; Brau et al. 2008), i.e., one can find explicit expressions for the particle densities $\sigma_0[E_0]$, $\rho_0[E_0]$ and for the front coordinate $\xi[E_0]$ as a function of the electric field E_0 . For the negative fronts treated here, the front is continuous as a function of D and the limit $D \rightarrow 0$ exists and equals the value of the front at $D = 0$, while for positive fronts ($E_\infty > 0$), it is singular (Ebert et al. 1997).

2.4 Multiplicity of Front Solutions, Pulled Fronts and Dynamical Selection

The nonionized state $(\sigma, \rho, E) = (0, 0, E_\infty)$ with a nonvanishing electric field E_∞ is linearly unstable under the temporal dynamics of the PDE (2.1)–(2.3). In fact, this spatial region ahead of the front dominates the dynamics, see the discussion in Ebert et al. (1997), Ebert and van Saarloos (2000a). Therefore, for fixed E_∞ , there is a continuous family of uniformly translating solutions, parameterized by the velocity $v \geq v^*$ (Ebert et al. 1996, 1997; Ebert and van Saarloos 1998, 2000a), where

$$v^*(E_\infty) = |E_\infty| + 2\sqrt{Df(|E_\infty|)}. \quad (2.8)$$

The dynamics of uniformly translating fronts with velocity $v > v^*$ are dominated by a flat spatial profile in the leading edge of the front

$$\sigma_v(\xi) \stackrel{\xi \rightarrow \infty}{\sim} e^{-\lambda \xi} \quad \text{with } \lambda < \Lambda^* = \sqrt{\frac{f(|E_\infty|)}{D}}, \quad (2.9)$$

where velocity v and decay rate λ are related through

$$v(E_\infty, \lambda) = |E_\infty| + D\lambda + \frac{f(E_\infty)}{\lambda}, \quad (2.10)$$

and therefore $v(E_\infty, \lambda) > v^*(E_\infty) \equiv v(E_\infty, \Lambda^*)$ for $\lambda \neq \Lambda^*$. The spatial profile (2.9) with $\lambda < \Lambda^*$ cannot build up dynamically from some initial condition with larger λ ; and it will destabilize if perturbed with an initial condition with smaller λ , therefore such flat and fast fronts can be approached dynamically only by initial conditions with exactly the same profile (2.9) in the leading edge. For a thorough discussion of this dynamics, we refer to Ebert and van Saarloos (2000a).

In practice, the continuum approximation for the electron density breaks down for very small densities in the leading edge and the initial electron distribution satisfies a decay condition of the form

$$\lim_{\xi \rightarrow \infty} \sigma(x, y, \xi, t = 0) e^{\lambda \xi} = 0 \quad \text{for all } \lambda < \Lambda^*, \quad (2.11)$$

if the penetrated state is really non-ionized. Therefore the velocity v^* is called the “selected” one, because it is the generic attractor for most physical initial conditions. Mathematically speaking, the profile with velocity v^* (the selected front) is the only profile that can build up dynamically from steeper initial conditions.

Therefore the condition (2.11) on the spatial decay of the initial electron distribution excludes all front solutions with velocity $v > v^*$ as long time attractors of the dynamics. If the criterion (2.11) is satisfied, then the selected front with speed v^* is dynamically stable and is approached with the universal algebraic convergence rate in time (Ebert and van Saarloos 1998, 2000a)

$$v(t) = v^* - \frac{3}{2\Lambda^* t} + \mathcal{O}\left(\frac{1}{t^{3/2}}\right). \quad (2.12)$$

However, without the spatial decay condition on the initial condition, the selected front is formally not stable (although this is physically irrelevant). This will lead to specific problems and solutions in the transverse stability analysis presented in the next section.

The spatial profile of the electron distribution in the selected front is

$$\sigma_{v^*}(\xi) \stackrel{\xi \rightarrow \infty}{\sim} (a\xi + b)e^{-\Lambda^* \xi}, \quad a > 0. \quad (2.13)$$

To summarize, if the analysis is restricted to initial conditions with a sufficiently rapid spatial decay in the electron densities (2.11), then the fronts have only one free external parameter, namely the field E_∞ ; it determines the asymptotic front velocity (2.8) and profile (2.13) after sufficiently long times. Furthermore, the equivariance in the system gives that the position of the front and its electrostatic potential are free internal parameters.

2.5 Full Spatial Profiles of the Selected Pulled Planar Front

The spatial decay behind the front will be important in the analysis as well, therefore we recall the basic behavior. For $\xi \rightarrow -\infty$, the electron density approaches

$$\sigma_{v^*}(\xi) \stackrel{\xi \rightarrow -\infty}{=} \sigma^- + ce^{\lambda^-\xi}, \quad c > 0, \quad (2.14)$$

and the electric field decays with the same exponent $E(\xi) = -(c/\lambda^-)e^{\lambda^-\xi}$. For $D = 0$,

$$\sigma^-(E_\infty, D = 0) = \int_0^{|E_\infty|} \alpha(x) dx \quad (2.15)$$

was derived in Ebert et al. (1997). For $D > 0$, σ^- decreases by a correction of order of D , more precisely,

$$\sigma^-(E_\infty, D) = \sigma^-(E_\infty, 0) + \mathcal{O}(D), \quad \sigma^-(E_\infty, D > 0) < \sigma^-(E_\infty, 0) \quad (2.16)$$

was proved in the Appendix of Li et al. (2007). The eigenvalue λ^- is given by

$$\lambda^- = \frac{\sqrt{v^{*2} + 4D\sigma^-} - v^*}{2D}, \quad (2.17)$$

where both v^* and σ^- depend on E_∞ and D . For small D , λ^- can be approximated as

$$\lambda^- = \frac{\sigma^-}{v^*} + \mathcal{O}(D) = \int_0^{|E_\infty|} \frac{\alpha(x) dx}{|E_\infty|} + \mathcal{O}(\sqrt{D}). \quad (2.18)$$

As an illustration, the spatial profiles of electron and ion density and the electric field of the selected front solution for a range of fields E_∞ and diffusion constants D are plotted in Fig. 1.

3 Numerical Calculation of the Dispersion Relation

First we will introduce the transversal perturbation setting and discuss an apparent degeneracy of the dispersion relation. However, it turns out that the constraint on the spatial decay of the electron density “selects” a single dispersion relation for every far field E_∞ . This relation then is calculated numerically based on dynamical systems techniques involving intersections of stable and unstable manifolds. Results for different fields E_∞ and diffusion constants D are presented.

3.1 Linear Transversal Perturbations of Planar Fronts

Suppose that there is a linear transversal perturbation of the uniformly translating front

$$\sigma(x, y, \xi, t) = \sigma_0(\xi) + \delta\bar{\sigma}_1(x, y, \xi, t) + \mathcal{O}(\delta^2), \quad \xi = z - v^*t, \quad (3.1)$$




Fig. 1 The pulled planar front solutions on the *left* for varying $E_\infty = -1, -5$ and -10 and fixed $D = 0.1$, and on the *right* for fixed $E_\infty = -1$ and varying $D = 0.1, 0.01$ and 0 . The *upper panels* show scaled electron and ion densities $\sigma_0(\xi)/\sigma^-(E_\infty, D)$ and $\rho_0(\xi)/\sigma^-(E_\infty, D)$ and the *lower panels* show the corresponding scaled electric fields $E_0(\xi)/|E_\infty|$ as a function of the spatial coordinate ξ . The fronts are displayed in a staggered way. The normalization factors $\sigma^-(E_\infty, D)$ in the *upper panels* are $\sigma^-(-1, 0) = 0.149$, $\sigma^-(-1, 0.01) = 0.148$, $\sigma^-(-1, 0.1) = 0.144$, $\sigma^-(-5, 0.1) = 2.832$, $\sigma^-(-10, 0.1) = 7.169$

and similarly for ρ and ϕ . The linearized equation for $\bar{\sigma}_1$, $\bar{\rho}_1$ and $\bar{\phi}_1$ follows from (2.1)–(2.3). By decomposing the perturbations into Fourier modes in the transversal directions x and y , by using isotropy in the transversal (x, y) plane and by using a Laplace transformation in t , the ansatz

$$(\bar{\sigma}_1, \bar{\rho}_1, \bar{\phi}_1) = e^{ikx+st}(\sigma_k, \rho_k, \phi_k)(\xi) \quad (3.2)$$

can be used for each Fourier component. The resulting equation can be written as a linear first-order system of ODEs, using the extra variables $\tau_k = \partial_\xi \sigma_k$ and $E_k = -\partial_\xi \phi_k$. Introduce $\mathbf{w} = (\tau_k, \sigma_k, \rho_k, E_k, \phi_k)$ and the linear system is given by

$$\partial_\xi \mathbf{w} = \mathbf{M}(\xi; E_\infty, k, s)\mathbf{w},$$

$$\text{with } \mathbf{M} = \begin{pmatrix} -\frac{E_0+v^*}{D} & \frac{2\sigma_0-\rho_0-f_0+s+Dk^2}{D} & -\frac{\sigma_0}{D} & -\frac{\partial_\xi \sigma_0-\sigma_0 f'_0}{D} & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{f_0}{v^*} & \frac{s}{v^*} & \frac{\sigma_0 f'_0}{v^*} & 0 \\ 0 & -1 & 1 & 0 & -k^2 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}. \quad (3.3)$$

In the matrix M , the abbreviated notations $f_0 = f(|E_0|)$ and $f'_0 = \partial_\eta f(\eta)|_{\eta=|E_0|}$ are used. For the terms with f'_0 , we have used that $E_0 < 0$, hence $\frac{E_0}{|E_0|} = -1$.

As the matrix M depends on k^2 , but not on k itself, the matrix is invariant under the transformation $k \rightarrow -k$. Thus if $s(k) = s^*$, then also $s(-k) = s^*$ and vice versa. Therefore it is sufficient to determine the dispersion relation for $k > 0$ and this will imply the relation for $k < 0$. From now on, we will use the convention that $k > 0$. Note that the invariance implies only that the dispersion relation will be a function of $|k|$. As will be shown later, the dispersion relation is not an analytic function of k near $k = 0$ and its expansion near $k = 0$ is linear in $|k|$.

For future use, we remark that the linearization matrix M does not involve any ξ -dependent terms in the fourth and fifth row and implies that E_k and ϕ_k are related by $E'_k = -\phi_k$. Thus the E_k -component of any solution of the linearized system (3.3) can be expressed as an integral

$$E_k(\xi) = c_1 e^{k\xi} + c_2 e^{-k\xi} + \frac{1}{2} \int_{\xi_0}^{\xi} [e^{k(\xi-\eta)} + e^{-k(\xi-\eta)}][\rho_k(\eta) - \sigma_k(\eta)] d\eta, \quad (3.4)$$

where the constants c_1 and c_2 are determined by the value of E_k and ϕ_k at $\xi = \xi_0$.

3.2 Stable and Unstable Manifolds and Degeneracy of the Dispersion Relation

The linearized problem (3.3) is a spectral problem with the spectral parameters s and k . If the asymptotic matrices $M^\pm(E_\infty, k, s) = \lim_{\xi \rightarrow \pm\infty} M(\xi; E_\infty, k, s)$ exist and are hyperbolic (i.e., no eigenvalues on the imaginary axis), then the system (3.3) has a bounded solution if and only if the unstable manifold from $\xi = -\infty$ and the stable manifold from $\xi = \infty$ have a nontrivial intersection. So we will focus in this section on determining the stable and unstable manifolds.

The behavior of the unstable manifold at the back of the front is given by the asymptotic matrix

$$\begin{aligned} M_-(E_\infty, k, s) &= \lim_{\xi \rightarrow -\infty} M(\xi; E_\infty, k, s) \\ &= \begin{pmatrix} -\frac{v^*}{D} & \frac{\sigma^- + s + Dk^2}{D} & -\frac{\sigma^-}{D} & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{s}{v^*} & 0 & 0 \\ 0 & -1 & 1 & 0 & -k^2 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}. \end{aligned}$$

For $s > 0$ and $k \neq 0$, this matrix has two negative and three positive eigenvalues:

$$\pm k, \quad \frac{s}{v^*}, \quad \mu_\pm = -\frac{v^*}{2D} \pm \frac{\sqrt{v^{*2} + 4D(\sigma^- + s + Dk^2)}}{2D}. \quad (3.5)$$

Thus the unstable manifold is three-dimensional. We remark that $\mu_+^-(s=0=k)$ is identical to the spatial decay rate λ^- (2.17) behind the unperturbed front.

Finding the stable manifold ahead of the front is less straightforward. Normally the stable manifold ahead of the front would be characterized by the matrix $\lim_{\xi \rightarrow +\infty} \mathbf{M}(\xi; E_\infty, k, s)$. For $s > 0$ and $s + Dk^2 < f(E_\infty)$ this matrix exists and has two positive and three negative eigenvalues:

$$\pm k, \quad \frac{s}{v^*}, \quad -\Lambda^* \pm \sqrt{\frac{s + Dk^2}{D}} = \frac{-\sqrt{f(E_\infty)} \pm \sqrt{s + Dk^2}}{\sqrt{D}}. \quad (3.6)$$

Thus the stable manifold is three-dimensional and a dimension count gives that the intersection of stable and unstable manifold is generically one-dimensional. So for small values of s and k , a continuous family of eigenvalues seems to exist. This feature is related to the instability of the asymptotic state at $+\infty$, to the continuous family of uniformly translating solutions for all $v \geq v^*(E_\infty)$, and to the instability of fronts against perturbations with smaller spatial decay rate λ , as discussed in the previous section. The continuous family of eigenvalues s for fixed wave number k is eliminated by applying the analysis only to fronts with a sufficiently rapid spatial decay (2.11). This condition will be imposed in the definition of the stable manifold; it will make the spectrum discrete.

Define the scaled vector

$$\tilde{\mathbf{w}} = \mathbf{D}\mathbf{w}, \quad \mathbf{D} = \text{diag}(e^{(\Lambda^* - \beta)\xi}, e^{(\Lambda^* - \beta)\xi}, 1, 1, 1), \quad (3.7)$$

where $\beta \in (0, \Lambda^*)$ will be fixed later and depend on k and Λ^* . The freedom in the choice of β is reminiscent of the fact that the decay condition holds for any $\lambda < \Lambda^*$, but not for $\lambda = \Lambda^*$. The system for $\tilde{\mathbf{w}}$ is

$$\tilde{\mathbf{w}}_\xi = \tilde{\mathbf{M}}(\xi; E_\infty, k, s)\tilde{\mathbf{w}}, \quad (3.8)$$

with

$$\tilde{\mathbf{M}} = \mathbf{D} \cdot \mathbf{M} \cdot \mathbf{D}^{-1} + (\partial_\xi \mathbf{D}) \cdot \mathbf{D}^{-1}$$

$$= \begin{pmatrix} -\frac{E_0 + v^*}{D} + \Lambda^* - \beta & \frac{2\sigma_0 - \rho_0 - f_0 + s + Dk^2}{D} & -\frac{\sigma_0}{D} e^{(\Lambda^* - \beta)\xi} & -\frac{\partial_\xi \sigma_0 - \sigma_0 f'_0}{D} e^{(\Lambda^* - \beta)\xi} & 0 \\ 1 & \Lambda^* - \beta & 0 & 0 & 0 \\ 0 & -\frac{f_0}{v^*} e^{-(\Lambda^* - \beta)\xi} & \frac{s}{v^*} & -\frac{-\sigma_0 f'_0}{v^*} & 0 \\ 0 & -e^{-(\Lambda^* - \beta)\xi} & 1 & 0 & -k^2 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix}.$$

Note that if $\beta = 0$, then the asymptotic matrix ahead of the front (at $\xi = +\infty$) does not exist because $e^{\Lambda^* \xi} \sigma_0(\xi)$ grows linearly in ξ according to (2.13). To get an asymptotic matrix ahead of the front, it is necessary that $0 < \beta < \Lambda^*$. In this case, the asymptotic matrix is

$$\tilde{\mathbf{M}}_+(E_\infty, k, s) = \lim_{\xi \rightarrow \infty} \tilde{\mathbf{M}}(\xi; E_\infty, k, s)$$

$$= \begin{pmatrix} -\Lambda^* - \beta & \frac{-f_\infty + s + Dk^2}{D} & 0 & 0 & 0 \\ 1 & \Lambda^* - \beta & 0 & 0 & 0 \\ 0 & -\frac{f_\infty}{v^*} & \frac{s}{v^*} & 0 & 0 \\ 0 & 0 & 0 & 0 & -k^2 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix},$$

where $f_\infty = f(|E_\infty|)$. The matrix $\tilde{\mathbf{M}}_+$ has the eigenvalues

$$\pm k, \quad \frac{s}{v^*}, \quad \text{and} \quad \mu_\pm^\pm = -\beta \pm \sqrt{\frac{s + Dk^2}{D}}. \quad (3.9)$$

Hence for $s > 0$ and $0 < \beta < \min(\Lambda^*, k\sqrt{1 + s/(Dk^2)})$, there are two negative and three positive eigenvalues. Thus the stable manifold of (3.8) is two-dimensional.

For the original unscaled system (3.3) this means that only the two-dimensional submanifold given by \mathbf{D}^{-1} acting on the stable manifold of (3.8) is relevant for the transverse instability. This submanifold will be called the stable manifold of (3.3) from now on. With this convention, the dispersion relation is a well-defined curve $s(k)$ and the curve is such that at $s = s(k)$, the linearized system (3.3) has a bounded solution which satisfies the spatial decay condition (2.11). Note that for both asymptotic matrices $\tilde{\mathbf{M}}_+$ and \mathbf{M}_- , the eigenvalues $\pm k$ become a degenerate eigenvalue 0 at $k = 0$. This leads to square root singularities and it can be expected that the dispersion relation $s(k)$ will be a function of $\sqrt{k^2} = |k|$ for k is small. This will be confirmed in Sect. 5.

3.3 The Evans Function for the Transverse Stability Problem

The occurrence of an intersection of the stable and unstable manifolds will be measured with the Evans function. Our numerical method to determine the dispersion curve as an eigenvalue problem is based on a definition of the Evans function in an exterior algebra framework and uses similar ideas as in Allen and Bridges (2002), Brin (2001), Brin and Zumbrun (2002), Blyuss et al. (2003), Bridges et al. (2003), Derks and Gottwald (2005). The approach of following the stable/unstable manifolds at $\xi = \pm\infty$ with a standard shooting method and checking their intersection using the Evans function, works only if these manifolds are one-dimensional or have codimension one; in the present model, this is the case in the singular limit $D = 0$ and a shooting method was used in Arrayás and Ebert (2004). Otherwise, any integration scheme will inevitably just be attracted by the eigendirection corresponding to the most unstable (stable) eigenvalue. Exterior algebra can be used to avoid this problem for higher dimensional manifolds and to preserve the analytic properties of the Evans function. Recently, a different method to calculate the Evans function for higher dimensional manifolds has been proposed in Humpherys and Zumbrun (2006). This method uses a polar coordinate approach and looks like a more suitable method for very high-dimensional problems.

To calculate the evolution of the two-dimensional stable and three-dimensional unstable manifold in a reliable numerical way, we will use the exterior algebra spaces

$\bigwedge^2(\mathbb{C}^5)$ and $\bigwedge^3(\mathbb{C}^5)$, respectively. The advantage of these spaces is that in $\bigwedge^l(\mathbb{C}^n)$, an l -dimensional linear subspace of \mathbb{C}^n can be described as a one-dimensional object, being the l -wedge product of a basis of this space. Also, the differential equation on \mathbb{R}^5 (or \mathbb{C}^5) induces a differential equation on the spaces $\bigwedge^l(\mathbb{C}^5)$:

$$\mathbf{W}_\xi = \mathbf{M}^{(l)}(\xi; E_\infty, k, s)\mathbf{W}, \quad \mathbf{W} \in \bigwedge^l(\mathbb{C}^5). \quad (3.10)$$

Here the linear operator (matrix) $\mathbf{M}^{(l)}$ is defined on a decomposable l -form $\mathbf{w}_1 \wedge \cdots \wedge \mathbf{w}_l$, $\mathbf{w}_i \in \mathbb{C}^5$, as

$$\mathbf{M}^{(l)}(\mathbf{w}_1 \wedge \cdots \wedge \mathbf{w}_l) := (\mathbf{M}\mathbf{w}_1) \wedge \cdots \wedge \mathbf{w}_l + \cdots + \mathbf{w}_1 \wedge \cdots \wedge (\mathbf{M}\mathbf{w}_l), \quad (3.11)$$

and it extends by linearity to the nondecomposable elements in $\bigwedge^l(\mathbb{C}^5)$. General aspects of the numerical implementation of this theory can be found in Allen and Bridges (2002). The general form of the matrices $\mathbf{M}^{(2)}$ and $\mathbf{M}^{(3)}$ can be found in the Appendix.

To determine the three-dimensional unstable manifold for $\xi \in (-\infty, 0]$, we will use (3.10) with $l = 3$. Since the induced matrix $\mathbf{M}^{(3)}(\xi; E_\infty, k, s)$ inherits the differentiability and analyticity of $\mathbf{M}(\xi; E_\infty, k, s)$, the following limiting matrix exists:

$$\mathbf{M}_-^{(3)}(E_\infty, k, s) = \lim_{\xi \rightarrow -\infty} \mathbf{M}^{(3)}(\xi; E_\infty, k, s).$$

The set of eigenvalues of the matrix $\mathbf{M}_\pm^{(3)}(E_\infty, k, s)$ consists of all possible sums of three eigenvalues of $\mathbf{M}_\pm(E_\infty, k, s)$ (see Marcus 1975). Therefore, for $s > 0$ and $k \neq 0$, there is an eigenvalue ν_- of $\mathbf{M}_-^{(3)}$, which is the sum of the three positive eigenvalues of \mathbf{M}_- , i.e.,

$$\nu_- = k + \frac{s}{v^*} - \frac{v^*}{2D} + \frac{\sqrt{v^{*2} + 4D(\sigma^- + s + Dk^2)}}{2D}$$

(note that the subscript “ $-$ ” in ν_- refers to exponentially decaying behavior at $-\infty$, not to the sign of ν_- , which is obviously positive). The eigenvalue ν_- is simple and has real part strictly greater than any other eigenvalue of $\mathbf{M}_-^{(3)}$ (as \mathbf{M}_- is hyperbolic). We denote the eigenvector associated with ν_- as \mathbf{W}_e^- , i.e., $\mathbf{M}_-^{(3)}\mathbf{W}_e^- = \nu_- \mathbf{W}_e^-$. This vector can always be constructed in an analytic way (see Kato 1984, pp. 99–101; Bridges et al. 2003; Brin and Zumbrun 2002; Humpherys et al. 2006). In this case it is easy to determine an explicit analytical expression for the eigenvector as \mathbf{M}_- is quite sparse. The unstable manifold corresponds to the solution $\mathbf{W}^-(\xi)$ of the linearized system (3.10) (with $l = 3$) which satisfies $\lim_{\xi \rightarrow -\infty} e^{-\nu_- \xi} \mathbf{W}^-(\xi) = \mathbf{W}_e^-$.

The stable manifold can be determined in a similar way. As indicated in the previous section, the scaled system (3.8) will be used to determine the stable manifold. For the stable manifold with $\xi \in [0, \infty)$, we will use (3.10) with $l = 2$ and the scaled matrix $\tilde{\mathbf{M}}$. As before, the asymptotic matrix

$$\mathbf{M}_+^{(2)}(E_\infty, k, s) = \lim_{\xi \rightarrow \infty} \tilde{\mathbf{M}}^{(2)}(\xi; E_\infty, k, s)$$

exists. Now the eigenvalues of $\mathbf{M}_+^{(2)}(E_\infty, k, s)$ consist of all possible sums of two eigenvalues of $\tilde{\mathbf{M}}_\pm(E_\infty, k, s)$. Therefore, for $s > 0, k \neq 0$, $\mathbf{M}_+^{(2)}$ has an eigenvalue ν_+ , which is the sum of the two negative eigenvalues of $\tilde{\mathbf{M}}_+$, i.e.,

$$\nu_+ = - \left(\sqrt{\frac{s + Dk^2}{D}} + k - \beta \right).$$

As before, this eigenvalue is simple and has real part strictly less than any other eigenvalue of $\mathbf{M}_+^{(2)}$. The eigenvector associated with ν_+ will be denoted by \mathbf{W}_e^+ , i.e., $\mathbf{M}_+^{(2)}\mathbf{W}_e^+ = \nu_+\mathbf{W}_e^+$. The stable manifold of the scaled system (3.8) corresponds to the solution $\mathbf{W}^+(\xi)$ of the linearized system (3.10) (with $l = 2$ and $\mathbf{M} = \tilde{\mathbf{M}}$) which satisfies $\lim_{\xi \rightarrow \infty} e^{-\nu_+\xi} \mathbf{W}^+(\xi) = \mathbf{W}_e^+$. To get the stable manifold of the original unscaled system, the inverse scalings matrix $\mathbf{D}^{-1}(\xi)$ has to be used. For arbitrary $\xi \geq 0$, the transformation in the wedge space $\bigwedge^2(\mathbb{C}^5)$ is quite complicated, but we will only need the original stable manifold at $\xi = 0$. And at $\xi = 0$, the scalings matrix is the identity matrix. Hence at $\xi = 0$, the scaled stable manifold and the original stable manifold are the same and $\mathbf{W}_e^+(0)$ describes the stable manifold of (3.3) at $\xi = 0$.

With the stable and unstable manifold as found above, the Evans function can be defined as

$$\Delta(E_\infty, k, s) = \mathbf{W}^-(0; E_\infty, k, s) \wedge \mathbf{W}^+(0; E_\infty, k, s), \quad s > 0, k \neq 0. \quad (3.12)$$

Thus the Evans function Δ is more or less the determinant of the matrix formed by a basis of the unstable manifold at $\xi = 0$ and a basis of the stable manifold at $\xi = 0$. If this function is zero, then the bases are linearly dependent, hence the two manifolds have a nontrivial intersection.

We have focused on the case $s > 0$. For $-Dk^2 < s < 0$, the system is still hyperbolic, but with a two-dimensional unstable manifold and a three-dimensional stable manifold. The above method can be easily adapted to calculate the dispersion curve in this region too.

3.4 Numerical Results on the Dispersion Relation with the Evans Function

To calculate the Evans function numerically, first the front solution has to be determined numerically as it appears explicitly in the linearization matrix $\mathbf{M}(\xi; E_\infty, k, s)$. The front is an invariant manifold connecting two fixed points of the ODE (2.6), so it can be easily determined by invariant manifold techniques or shooting, using the package DStool (Back et al. 1992). Shooting works in this case as the front connects a one-dimensional unstable manifold to a three-dimensional center-stable manifold in the ODE (2.6).

After determining the fronts, the stable and unstable manifolds can be calculated by numerical integration, see e.g. Allen and Bridges (2002), Bridges et al. (2003), Brin and Zumbun (2002). In the numerical calculation of the stable manifold, we will use $\beta = \frac{1}{2} \min(\Lambda^*, k)$. For the stable manifold, the linearized equation on $\bigwedge^2(\mathbb{C}^5)$

$$\widehat{\mathbf{W}}_\xi^+ = [\tilde{\mathbf{M}}^{(2)}(\xi; E_\infty, k, s) - \nu_+(E_\infty, k, s)\mathbf{I}]\widehat{\mathbf{W}}^+, \quad \widehat{\mathbf{W}}^+(\xi)|_{\xi=L_\infty} = \mathbf{W}_e^+(E_\infty, k, s)$$

is integrated from $x = L_\infty$ to $\xi = 0$, using the second-order Gauss–Legendre Runge–Kutta (GLRK) method, i.e., the implicit midpoint rule. Here the scaling $\widehat{\mathbf{W}}^+(\xi) = e^{-\nu_+\xi} \mathbf{W}^+(\xi)$ ensures that any numerical errors due to the exponential growth are removed and $\widehat{\mathbf{W}}^+(\xi)|_{\xi=0} = \mathbf{W}^+(\xi)|_{\xi=0}$ is bounded. The eigenvector $\mathbf{W}_e^+(E_\infty, k, s)$ can be determined explicitly as wedge product of the relevant eigenvectors of $\mathbf{M}^+(E_\infty, s, k)$ thanks to the sparse nature of this matrix.

For the unstable manifold, the linearized equation on $\bigwedge^3(\mathbb{C}^5)$

$$\widehat{\mathbf{W}}_\xi^- = [\mathbf{M}^{(3)}(\xi; E_\infty, k, s) - \nu_-(E_\infty, s, k)\mathbf{I}]\widehat{\mathbf{W}}^-, \quad \widehat{\mathbf{W}}^-(\xi)|_{\xi=L_\infty} = \mathbf{W}_e^-(E_\infty, s, k)$$

is integrated from $x = -L_\infty$ to $\xi = 0$, also using the implicit midpoint rule and introducing the rescaling $\widehat{\mathbf{W}}^-(\xi) = e^{-\nu_-\xi} \mathbf{W}^-(\xi)$ to remove potential exponential growth. Again, the eigenvector $\mathbf{W}_e^-(E_\infty, k, s)$ can be determined explicitly as wedge product of the relevant eigenvectors of $\mathbf{M}^-(E_\infty, s, k)$.

At $\xi = 0$, the computed Evans function is (see (3.12))

$$\Delta(E_\infty, k, s) = \mathbf{W}^-(0) \wedge \mathbf{W}^+(0) = \widehat{\mathbf{W}}^-(0) \wedge \widehat{\mathbf{W}}^+(0). \quad (3.13)$$

For $s = 0 = k$, the center-stable and the center-unstable manifold have a two-dimensional intersection, due to the translation and gauge invariance, see Sect. 5.1 for details. In order to determine the dispersion curve, we start near $k = 0$ and $s = 0$ and then slowly increase k and determine for which $s(k)$ the Evans function $\Delta(E_\infty, k, s(k))$ vanishes.


The numerical errors in the calculation of the Evans function are mainly influenced by the step size used in the numerical integration with the GLRK method and errors in the numerically determined front. The numerical integration uses the step size $\delta x = 0.01$. We have performed various checks with a decreased step size, and these checks show that the error in the value of s for fixed k is largest (order 10^{-4}) if k is small and decreases for larger k (order 10^{-6}). The accuracy of the front has been checked and is such that the error in the front gives a negligible error (compared to the error due to the error in the step size) in the value of $s(k)$. It turns out that the scheme is not very sensitive to errors in the front (at least for the E_∞ and D values considered).

In the following sections, we present data for the dispersion curve for the varying electric field E_∞ and diffusion coefficient D . A more detailed discussion of the data, its relation with analytical asymptotics and some empirical fitting can be found in Sect. 6.


3.4.1 Varying the Electric Field ahead of the Front

First we consider how the dispersion curve depends on the electric field E_∞ ahead of the front, while the diffusion coefficient is fixed to $D = 0.1$. In Fig. 2a, the dispersion curve is shown for $E_\infty = -1$, $E_\infty = -5$ and $E_\infty = -10$. The figure shows that the shape of the dispersion curve stays similar, but the scales of s and k increase when E_∞ increases. The dispersion curves can be characterized by the maximal growth rate s_{\max} and the corresponding wave number k_{\max} where $s(k_{\max}) = s_{\max}$ as well as by the wave number $k_0 > 0$ with $s(k_0) = 0$ that limits the band $0 < k < k_0$ of wave numbers with positive growth rates.

Fig. 2 Dispersion curves $s(k)$: (a) for varying E_∞ and fixed $D = 0.1$, and (b) for fixed $E_\infty = -1$ and varying D . The pairs (E_∞, D) shown are the same as in Fig. 1. The data for the singular limit $D = 0$ are taken from Arrayás and Ebert (2004)



(a) $E_\infty = -1, -5$ and -10 and fixed $D = 0.1$



(b) Fixed $E_\infty = -1$ and $D = 0.1, 0.01$ and 0

3.4.2 Varying the Diffusion Coefficient

Next we consider the effect of varying the diffusion coefficient D , while keeping the electric field ahead of the front fixed at $E_\infty = -1$. In Arrayás and Ebert (2004) it is shown that if diffusion is ignored ($D = 0$), the dispersion curve stays positive and is monotonically increasing to the saturation value $s(k) = f(|E_\infty|)/2$ for $k \rightarrow \infty$. Our numerics show that if diffusion is present, this is not the case anymore. This is not surprising as diffusion is a singular perturbation. In Fig. 2b, the dispersion curve is shown for $D = 0.1$, $D = 0.01$ and $D = 0$; the data for $D = 0$ are taken from Arrayás and Ebert (2004). It shows that the growth rate $s(k)$ has a maximum s_{\max} if diffusion is present and becomes negative for k larger than some k_0 . Furthermore, for decreasing diffusion D , the maximal growth rate moves upward toward the saturation value $f(|E_\infty|)/2$ for $D = 0$. This suggests that some features of the dispersion curve behave regularly in D , in spite of the fact that D is a singular perturbation. For example, for a finite wave number interval, the limit of the dispersion curves for $D \rightarrow 0$ exists and is the curve for $D = 0$. However, the asymptotic profile for large values of the

wave number is obviously singular in D . This duality can also be found in the front itself: the velocity and the profile of the ionization density and the electric field of the uniformly translating negative front depend regularly on $D = 0$, while the profile of the ionization density is singular, as discussed in Sect. 2.4 and shown in Fig. 1.

4 Numerical Simulation of the Perturbed Initial Value Problem

In the previous section, we determined the dispersion relation $s(k)$ for transversal perturbations of ionization fronts as a temporal eigenvalue problem of the PDE system linearized about the uniformly translating planar front. Since we are dealing with pulled fronts (cf. Sects. 1 and 2.4), the problem is unconventional: both the velocity v^* of the uniformly translating planar front and the dispersion relation $s(k)$ of its transversal perturbations are unique only if the spatial decay constraint (2.11) is imposed. Furthermore a longitudinally perturbed planar front approaches its asymptotic profile and velocity algebraically slowly in time (2.12). Therefore it is worthwhile to test the predicted dispersion relation on direct numerical simulations of the corresponding initial value problem.

In this section, we will therefore simulate the temporal evolution of a perturbed planar front by numerically solving the full nonlinear PDEs (2.1)–(2.3), and we will determine the dispersion curve from a number of simulations with perturbations with different wave vectors k . This is done for far field $E_\infty = -1$ and diffusion constant $D = 0.1$.

To determine the instability curve with a simulation of the full PDE, we parameterize the evolution of a perturbed planar front with wave number k as

$$\mathbf{U}(x, z, t) \approx \mathbf{U}_0(\xi) + \delta \mathbf{U}_1(\xi, t) e^{ikx+st}, \quad \xi = z - v^*t, \quad \mathbf{U} = (\sigma, \rho, \phi). \quad (4.1)$$

If δe^{st} is small enough, the solution is in the linear regime, and $s(k)$ can be determined from the evolution of the perturbation after $\mathbf{U}_1(\xi, t)$ has relaxed to some time independent function. Therefore in the numerical simulations, we choose δ for each wave number k in such a manner that t is large enough to extract meaningful growth rates and that δe^{st} is small enough that the dynamics at the final time is still well approximated by the linearization about the planar front.

Furthermore, an appropriate choice of the initial condition reduces the initial transient time during which $\mathbf{U}_1(\xi, t)$ in the comoving frame still explicitly depends on time t . Ideally, such an initial condition is of the form $\mathbf{U}(x, z, 0) = \mathbf{U}_0(\xi) + \delta \mathbf{U}_1(\xi) \cos kx$ etc., where \mathbf{U}_1 is a solution of the linearized system (3.3). To find an approximation for $\mathbf{U}_1(\xi)$, we use that the instability acts on the position of the front, i.e., we write the perturbed front as $\mathbf{U}_0(\xi + \delta e^{ikx+st}) \approx \mathbf{U}_0(\xi) + \delta e^{ikx+st} \partial_\xi \mathbf{U}_0(\xi)$. Therefore we choose $\mathbf{U}_1(\xi) = \partial_\xi \mathbf{U}_0(\xi)$ and the initial condition as

$$\mathbf{U}(x, z, 0) = \mathbf{U}_0(z) + \delta \partial_z \mathbf{U}_0(z) \cos kx. \quad (4.2)$$

As $\partial_\xi \mathbf{U}_0(\xi)$ is a solution of the linearized system for $k = 0 = s$, this choice will be very efficient for small values of k and require longer transient times for larger k .

To solve the full 2D PDE, the algorithm as described in Arrayás et al. (2002), Rocco et al. (2002) is used, while adaptive grid refinement as introduced in Montijn et al. (2006b) is not required. For fixed k , the PDE with initial condition (4.2) is solved on the spatial rectangle $(x, z) \in [0, L_x] \times [0, L_z]$. The length of the domain in the transversal x -direction, L_x , is such that exactly five wavelengths fit into the domain, i.e., $L_x = \frac{10\pi}{k}$, and periodic boundary conditions are imposed in this direction by identifying $x = 0$ with $x = L_x$. On the boundaries in the longitudinal z -direction, Neumann conditions for the electron density are imposed. The potential is constant behind the front and the electric field is constant ahead of the front; therefore for the potential ϕ , the Dirichlet condition $\phi = 0$ is imposed at $z = 0$, and the Neumann condition $\partial_z \phi = -E_\infty$ at $z = L_z$ accounts for the far field ahead of the front.

The amplitude of the perturbation is conveniently traced by the maximum of the electron density

$$\sigma_{\max}(x, t) = \max_{z \in [0, L_z]} \sigma(x, z, t) \quad (4.3)$$

evaluated across the front. The reason is as follows. First, Fig. 1 shows the spatial profiles of planar fronts for different electric fields E_∞ and illustrates that, for fixed D , the maximum of the electron density σ_{\max} as well as the asymptotic density σ^- behind the front strongly depend on the field E^+ immediately ahead of the front; see also (2.15) and (2.16). Here we consider a planar front, thus the close and the far field are identical: $E^+ = E_\infty$. Second, the modulation of the front position leads to a modulation of the electric field E^+ immediately before the front (cf. discussion in Sect. 5.2); therefore σ_{\max} as a function of E^+ is modulated as well.

An example of $\sigma_{\max}(x, t)$ as a function of the transversal coordinate x for a fixed time t is plotted in Fig. 3a. The amplitude of the wave modulation is determined by the Fourier integral


$$A(t, k) = \frac{k}{5\pi} \int_0^{\frac{10\pi}{k}} \sigma_{\max}(x, t) \cos kx \, dx.$$

In Fig. 3b we plot $\log A$ against time t for $k = 0.45$. Note that $k = 0.45$ is close to $k_0 = 0.482$ (see Fig. 2a and Table 1) where the growth rate vanishes, $s(k_0) = 0$, therefore the growth rate in the present example is small and particularly sensitive to numerical errors.


Figure 3b shows an initial temporal transient before steady exponential growth is reached (where exponential growth manifests itself as a straight line in the logarithmic plot). This is typically observed for the larger k -values ($k > 0.1$); as said before, this is related to the fact that the function $\mathbf{U}_1(z)$ in the initial condition (4.2) is not optimal. For $k < 0.1$, there are less transients as $\mathbf{U}_1(z) \approx \partial_z \mathbf{U}_0(z)$ for small values of k .

To determine the growth rate $s(k)$, a least squares algorithm is used to fit the best line through the data points $(t, \log A)$, and the initial transient time is ignored for larger values of k . For each value of k , the growth rate is determined with various choices of δ . The resulting growth rate $s(k)$ is indicated in Fig. 4 with crosses \times and the error bars are related to the various choices of δ .

Fig. 3 Examples of data of the initial value simulation from which the growth rate $s(k)$ shown in Fig. 4 are determined



(a) The maximal value of the electron density $\sigma_{\max}(x, t)$ for $t = 50$ as a function of the transversal coordinate x . The perturbation has wave number $k = 0.45$, the transversal length $L_x = 10\pi/k$ leaves space for five wavelengths that are clearly visible



(b) The logarithm of the amplitude of the front modulation $\log A$ as a function of time t for the same k

Figure 4 also shows the dispersion relation for $(E_\infty, D) = (-1, 0.1)$ determined with the dynamical systems method in the previous Sect. 3.4; these numerical results are denoted with $+$ and can now be compared with the results of the initial value problem from the present section. Around the maximum of the curve, the agreement between the numerical results of the two very distinct methods is convincing. For larger values of k , the differences increase, but the error bars of the initial value problem results increase as well. Furthermore, the plotted error bars are an underestimation, as they only account for the errors discussed earlier. These errors emerge from the choice of the initial condition and from the time interval of evaluation and




Fig. 4 The dispersion curve $s(k)$ for $E_\infty = -1$ and $D = 0.1$. The crosses \times with error bars indicate results of simulations of the full initial value problem as discussed in Sect. 4 and demonstrated in Fig. 3a. For comparison, the results of the dynamical systems method from Sect. 3.4 are indicated with $+$ symbols

therefore from possible initial transients and from a possible transition to nonlinear behavior. Additional errors can be due to the numerical discretization and time stepping of the PDEs themselves. We therefore conclude that the two results agree within the numerical error range of the initial value simulations over the whole curve.

5 Analytical Derivation of Asymptotic Limits for $k \ll 1$ and $k \gg 1$

Having determined the dispersion relation numerically for different values of electric field E_∞ and diffusion constant D in Sect. 3, and having tested the correctness of the eigenvalue calculation against numerical solutions of the initial value problem in Sect. 4, we now analytically derive asymptotic expressions for the dispersion relation for small and large values of the wave modes k . It will be shown that these asymptotic limits are

$$s(k) = \begin{cases} kE_\infty \frac{dv^*}{dE_\infty}, & k \ll 1, \\ -Dk^2, & k \gg 1. \end{cases}$$

In doing so, we mathematically formalize and generalize the derivation of the small k asymptotic limit that was presented in Arrayás and Ebert (2004) for the singular limit $D = 0$, and we correct the result proposed in Arrayás et al. (2005); we also rigorously derive the large k asymptotic limit, in agreement with the form proposed in Arrayás et al. (2005).

5.1 Analysis for the Asymptotic Limit $k \ll 1$

Translation invariance and electrostatic gauge invariance give two explicitly known bounded solutions of the linearized system (3.3) at $k = 0$ and $s = 0$. These are

$$\mathbf{u}'_0(\xi) = (\sigma''_0(\xi), \sigma'_0(\xi), \rho'_0(\xi), E'_0(\xi), -E_0(\xi))^T \quad \text{and} \quad \mathbf{e}_5 = (0, 0, 0, 0, 1)^T.$$

Note that \mathbf{e}_5 is a solution of the linearized system (3.3) for $k = 0$ and s arbitrary.

From the asymptotics of (3.3) for $k = 0 = s$ at $\xi = -\infty$, we see that the only exponentially decaying solution at $\xi = -\infty$ is given by $\mathbf{u}'_0(\xi)$. This solution is related to the only positive eigenvalue μ_+^- (see (3.5)). For $\xi \rightarrow +\infty$, the solution $\mathbf{u}'_0(\xi) \rightarrow -E_\infty \mathbf{e}_5$, hence this solution is not exponentially decaying for $\xi = +\infty$. However, it is easy to obtain an explicit exponentially decaying solution at $\xi = +\infty$, this is $\mathbf{u}'_0(\xi) + E_\infty \mathbf{e}_5$.

From the eigenvalues in (3.5) it follows that for $0 < k \ll 1$ and $0 < s \ll 1$, the three-dimensional unstable manifold at $\xi \rightarrow -\infty$ involves one eigenfunction with a fast exponential decay (related to the eigenvalue μ_+^-) and two eigenfunctions with a slow exponential decay (related to the eigenvalues k and $\frac{s}{v^*}$). Similarly, from the eigenvalues in (3.9), it follows that the two-dimensional stable manifold at $\xi \rightarrow +\infty$ involves one eigenfunction with a fast exponential decay (related to the eigenvalue $-\Lambda^* + \beta + \mu_+^+$) and one eigenfunction with a slow exponential decay (related to the eigenvalue $-k$). Recall that the stable manifold is defined as a subset of the full stable manifold to account for the spatial decay condition (2.11).

We focus on approximating an exponentially decaying solution on the stable manifold. As we have seen above, in lowest order, this solution is

$$\mathbf{w}^s(\xi; E_\infty, 0, 0) = \mathbf{u}'_0(\xi) + E_\infty \mathbf{e}_5 + \mathcal{O}(k + s).$$

To determine the next order, we will use the slow behavior of the asymptotic system and write

$$\begin{aligned} \mathbf{w}^s(\xi; E_\infty, k, s) &= \mathbf{u}'_0(\xi) + E_\infty(0, 0, 0, k, 1)e^{-k\xi} + kU_{1,k}^s(\xi) \\ &\quad + sU_{1,s}^s(\xi) + \mathcal{O}((s + k)^2). \end{aligned}$$

The second term on the right-hand side ($E_\infty(0, 0, 0, k, 1)e^{-k\xi}$) is an approximation of the slow behavior on the stable manifold, while the other three terms are related to the fast decay. Because of the slow decay, the expansion is only valid on a ξ -interval with $k\xi = o(1)$, hence ξ should not be too large.

Substitution of these expressions into the linearized system (3.3) gives that $U_{1,s}^s$ and $U_{1,k}^s$ have to satisfy

$$\begin{aligned} (D_\xi - \mathbf{M}(\xi; E_\infty, 0, 0))U_{1,s}^s &= \left(\frac{\sigma'_0}{D}, 0, \frac{\rho'_0}{v^*}, 0, 0 \right), \\ (D_\xi - \mathbf{M}(\xi; E_\infty, 0, 0))U_{1,k}^s &= -E_\infty \left(-\frac{\tau_0 - \sigma_0 f'_0}{D} e^{-k\xi}, 0, \frac{\sigma_0 f'_0}{v^*} e^{-k\xi}, 0, 1 - e^{-k\xi} \right). \end{aligned}$$

By analyzing the unperturbed system, we can find a particular solution of the first equation. The front solution $\mathbf{u}_0 = (\tau_0, \sigma_0, \rho_0, E_0, \phi_0)$ satisfies

$$\begin{aligned}\tau'_0 &= -\frac{v^* + E_0}{D}\tau_0 + (\sigma_0 - \rho_0)\frac{\sigma_0}{D} - \frac{\sigma_0 f_0}{D}, \\ \sigma'_0 &= \tau_0, \\ \rho'_0 &= -\frac{\sigma_0 f_0}{v^*}, \\ E'_0 &= -(\sigma_0 - \rho_0), \\ \phi'_0 &= -E_0.\end{aligned}$$

Differentiating this system with respect to E_∞ gives

$$\begin{aligned}(D_\xi - \mathbf{M}(\xi; E_\infty, 0, 0))\frac{\partial \mathbf{u}_0}{\partial E_\infty} &= \frac{dv^*}{dE_\infty} \left(-\frac{\tau_0}{D}, 0, \frac{\sigma_0 f_0}{(v^*)^2}, 0, 0 \right) \\ &= -\frac{dv^*}{dE_\infty} \left(\frac{\sigma'_0}{D}, 0, \frac{\rho'_0}{v^*}, 0, 0 \right).\end{aligned}$$

Hence

$$U_{1,s}^s = -\left(\frac{dv^*}{dE_\infty}\right)^{-1} \frac{\partial \mathbf{u}_0}{\partial E_\infty} + \text{a homogeneous solution.}$$

The asymptotic behavior of $\frac{\partial \mathbf{u}_0}{\partial E_\infty}$ is

$$\frac{\partial \mathbf{u}_0}{\partial E_\infty} \sim (0, 0, 0, 1, -\xi), \quad \xi \rightarrow \infty. \quad (5.1)$$

The polynomial growth in the ϕ_k -component will need to be canceled by the behavior of the other terms which involve k and hence will give a relation between s and k .

In fact, the E_k -components of the linearized system (3.3) for any s or k can be expressed by the integral equation (3.4). For all solutions on the stable manifold, the limit $E_k(\infty) = \lim_{\xi \rightarrow \infty} E_k(\xi)$ is well defined, so we can write the E_k -components on the stable manifold as

$$E_k^s(\xi) = c_2 e^{-k\xi} - \frac{1}{2} \int_\xi^\infty [e^{k(\xi-\eta)} + e^{-k(\xi-\eta)}][\rho_k(\eta) - \sigma_k(\eta)] d\eta.$$

In the integral, σ_k must satisfy the decay condition (2.11) on the stable manifold and hence will have fast exponential decay. Furthermore, from (3.3), it can be seen that $\rho_k(\xi) = c_3 e^{\frac{s\xi}{v^*}} + \frac{1}{v^*} \int_\xi^\infty e^{-\frac{s(\xi-\eta)}{v^*}} [\sigma_0(\eta) f'_0(\eta) E_k(\eta) - f_0(\eta) \sigma_k(\eta)] d\eta$. As the term inside the integral has fast exponential decay, we get that on the stable manifold $c_3 = 0$ and ρ_k has fast exponential decay too. Thus the integral in the expression for E_k^s has fast exponential decay for large ξ . Since $\phi_k = -E'_k$, we get on the stable manifold for small k and ξ -values not too large, say $\xi \sim k^{-\frac{1}{4}}$ (hence $k\xi \sim k^{\frac{3}{4}}$)

$$\begin{aligned}(E_k^s, \phi_k^s)(\xi) &= (k, 1)e^{-k\xi} + \mathcal{O}(e^{-\Lambda^* \xi}) \\ &= (k, 1)(1 - k\xi + \mathcal{O}(k\sqrt{k})) = (k, 1 - k\xi) + \mathcal{O}(k\sqrt{k}).\end{aligned}$$

The exponentially decaying solution on the stable manifold is given by

$$\begin{aligned} \mathbf{w}^s(\xi; E_\infty, k, s) = & \mathbf{u}'_0(\xi) - s \left(\frac{dv^*}{dE_\infty} \right)^{-1} \frac{\partial \mathbf{u}_0}{\partial E_\infty}(\xi) + E_\infty(0, 0, 0, k, 1)e^{-k\xi} \\ & + kU_{1,k}^s(\xi) + \mathcal{O}((s+k)^2), \end{aligned}$$

and the arguments above show that the order k contribution in the (E_k, ϕ_k) -components is given fully by $E_\infty(0, 0, 0, k, 1)e^{-k\xi}$ and that $kU_{1,k}^s(\xi)$ does not contribute to those components at this order. So it follows that the polynomial growth in the ϕ_k -component of $\frac{\partial \mathbf{u}_0}{\partial E_\infty}(\xi)$ as given by (5.1) has to be canceled by the ϕ_k -component in $E_\infty(0, 0, 0, k, 1)e^{-k\xi}$, i.e., $s(\frac{dv^*}{dE_\infty})^{-1} = kE_\infty$ or

$$s = c^*k + \mathcal{O}(k^2), \quad \text{with } c^* = E_\infty \frac{dv^*}{dE_\infty}, \quad (5.2)$$

and v^* given in (2.8). Equation (5.2) establishes the small k limit of the dispersion relation $s(k)$.

5.2 A Physical Argument for the $k \ll 1$ Asymptotic Limit

There is also a physical argument for the asymptotic limit (5.2) that generalizes the calculation in Arrayás and Ebert (2004, Sect. IV.C) to nonvanishing $D > 0$. For $k \ll 1$, the wavelength of the transversal perturbation $2\pi/k$ is the largest length scale of the problem. It is much larger than the inner longitudinal structure of the ionization front. On the length scale $2\pi/k$, the front can therefore be approximated by a moving boundary between the ionized and nonionized regions at the position

$$z_f(x, t) = z_0 + v^*(E_\infty)t + \delta e^{ikx+st}, \quad (5.3)$$

and the local velocity of this perturbed front is

$$v(x, t) = \partial_t z_f(x, t) = v^*(E_\infty) + s\delta e^{ikx+st}. \quad (5.4)$$

The electric field in the nonionized region is determined by $\mathbf{E} = -\nabla\phi$, where ϕ is the solution of the Laplace equation $\nabla^2\phi = 0$ together with the boundary conditions; these are $\mathbf{E} \rightarrow E_\infty\hat{\mathbf{z}}$ for $z \rightarrow \infty$ fixing the field far ahead of the front and $\phi(x, z_f, t) = 0$ making the ionization front equipotential. (Due to gauge invariance the constant potential can be set to zero.) The solution of this problem is

$$\begin{aligned} \phi(x, z, t) = & -E_\infty(z - z_0 - v^*t) \\ & + E_\infty e^{-k(z-z_f)}\delta e^{ikx+st} + \mathcal{O}(\delta^2), \quad \text{for } z \geq z_f, \\ E^+(x, t) = & E_\infty + kE_\infty\delta e^{ikx+st} + \mathcal{O}(\delta^2), \quad \text{for } z = z_f, \end{aligned} \quad (5.5)$$

here $E^+(x, t) = \lim_{\epsilon \downarrow 0} E(x, z_f + \epsilon, t)$ is the electric field extrapolated onto the boundary from the nonionized side.

As the perturbation is linear, the front is almost planar $\delta \ll 2\pi/k$. Therefore it will propagate with the velocity $v^*(E^+) = |E^+| + 2\sqrt{Df(|E^+|)}$ (2.8) of the planar front in the local field E^+ . Inserting E^+ from (5.5) and expanding about E_∞ , we get

$$v(x, t) = v^*(E^+(x, t)) = v^*(E_\infty) + \partial_E v^*|_{E_\infty} k E_\infty \delta e^{ikx+st} + O(\delta^2). \quad (5.6)$$

Comparison of (5.6) and (5.4) immediately gives the dispersion relation $s = c^*k + O(k^2)$ (5.2), that generalizes the result $s(k) = |E_\infty|k + O(k^2)$ that was derived in Arrayás and Ebert (2004) for the singular limit $D = 0$.

5.3 Analysis for the Asymptotic Limit $k \gg 1$

The asymptotic limit for $k \gg 1$ is derived by a contradiction argument. We will suppose that k is large and that $s + Dk^2$ is positive, but not small, i.e.,

$$k \gg 1, \quad s + Dk^2 > 0, \quad \text{and} \quad s + Dk^2 \neq o(1) \quad (5.7)$$

and show that this does not allow for bounded solutions. With the assumptions above on s and k , the dominant contributions in the matrix \mathbf{M} on the whole axis ξ are

$$\mathbf{M}_\infty = \begin{pmatrix} 0 & \frac{s+Dk^2}{D} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{s}{v^*} & 0 & 0 \\ 0 & 0 & 0 & 0 & -k^2 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix} + O(1). \quad (5.8)$$

Here the three entries -1 , 1 and $\frac{s}{v^*}$ are necessary for a nonvanishing determinant.

We want to use the Roughness Theorem (Coppel 1978) for exponential dichotomies to show that for k large and s not close to $-Dk^2$, the exponential dichotomy of the constant coefficient ODE is close to the exponential dichotomy of the full system. So first we recall the definition of an exponential dichotomy, which gives projections on stable or unstable manifolds.

Definition 1 (Coppel 1978) Let \mathbf{A} be a matrix in $\mathbb{R}^{n \times n}$, $u \in \mathbb{R}^n$, and $J = \mathbb{R}_-, \mathbb{R}_+$, or \mathbb{R} . Let $\Phi(y)$ be a solution matrix of the linear system

$$\frac{du}{dy} = \mathbf{A}(y)u, \quad y \in J. \quad (5.9)$$

The linear system (5.9) is said to possess an exponential dichotomy on the interval J if there exist a projection \mathbf{P} and constants K and $\kappa^s < 0 < \kappa^u$ with the following properties:

$$\begin{aligned} |\Phi(y)\mathbf{P}\Phi^{-1}(y_0)| &\leq K e^{\kappa^s(y-y_0)}, \quad \text{for } y \geq y_0, \quad y, y_0 \in J, \\ |\Phi(y)(\mathbf{I} - \mathbf{P})\Phi^{-1}(y_0)| &\leq K e^{\kappa^u(y-y_0)}, \quad \text{for } y_0 \geq y, \quad y, y_0 \in J. \end{aligned}$$

An extension for PDEs of this definition can be found in Peterhof et al. (1997).

The Roughness Theorem for exponential dichotomies states the following.

Theorem 2 (Roughness Theorem, Coppel 1978) *Consider the system*

$$\frac{du}{dy} = [\mathbf{A}_0 + \mathbf{A}_1(y)]u, \quad (5.10)$$

with $\mathbf{A}_0 \in \mathbb{R}^{n \times n}$ a hyperbolic matrix and $u \in \mathbb{R}^n$. Then for all $\delta_0 > 0$ there exists a $\delta_1 > 0$ such that for all matrix functions $\mathbf{A}_1 : \mathbb{R} \rightarrow \mathbb{R}^{n \times n}$ with $\|\mathbf{A}_1\|_{L^\infty(\mathbb{R}^+, \mathbb{R}^{n \times n})} < \delta_1$, the system (5.10) has an exponential dichotomy on \mathbb{R}^+ (and \mathbb{R}^-) with its dichotomy exponents and projections δ_0 -close to those of $\frac{du}{dy} = \mathbf{A}_0 u$ (in the $L^\infty(\mathbb{R}^+, \mathbb{R}^{n \times n})$ norm).

A constant coefficient linear system does not have bounded solutions. So if the exponential dichotomy of the linearized system (3.3) is close to the dichotomy of the constant coefficient system with the matrix \mathbf{M}_∞ as in (5.8), the linearized system (3.3) does not have bounded solutions either. We will show that this is the case if s and k satisfy the assumptions (5.7).

First, we introduce some scaling and coordinate transformations. Define the small parameter $\varepsilon = \frac{1}{k}$ and the scaled spatial variable, the transformation matrix and transformed vector

$$\eta = k\xi = \frac{\xi}{\varepsilon}, \quad \mathbf{T}(\varepsilon) = \text{diag}(\varepsilon, 1, \varepsilon, \varepsilon, 1) \quad \text{and} \quad \widehat{\mathbf{w}}(\eta) = \mathbf{T}(\varepsilon)\mathbf{w}(\varepsilon\eta).$$

Now (3.3) can be written as

$$\partial_\eta \widehat{\mathbf{w}} = [\widehat{\mathbf{M}}_0(\varepsilon, s) + \varepsilon \widehat{\mathbf{M}}_1(\eta; E_\infty, \varepsilon)] \widehat{\mathbf{w}}, \quad (5.11)$$

with

$$\widehat{\mathbf{M}}_0(\varepsilon, s) = \begin{pmatrix} 0 & 1 + \frac{\varepsilon^2 s}{D} & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{s\varepsilon}{v} & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & -1 & 0 \end{pmatrix},$$

and

$$\widehat{\mathbf{M}}_1(\eta; E_\infty, \varepsilon) = \begin{pmatrix} -\frac{E_0 + v^*}{D} & \varepsilon \frac{2\sigma_0 - \rho_0 - f_0}{D} & -\frac{\sigma_0}{D} & -\frac{\partial_\xi \sigma_0 - \sigma_0 f'_0}{D} & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -\varepsilon \frac{f_0}{v^*} & 0 & -\frac{\sigma_0 f'_0}{v^*} & 0 \\ 0 & -\varepsilon & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

where $\xi = \varepsilon\eta$.

The eigenvalues and eigenvectors of the constant coefficient matrix $\widehat{\mathbf{M}}_0$ are

$$\begin{aligned} \pm 1, & \quad \text{with eigenvectors } \mathbf{w}_{\pm 1} = (0, 0, 0, \mp 1, 1), \\ \pm \sqrt{1 + \frac{\varepsilon^2 s}{D}}, & \quad \text{with eigenvectors } \mathbf{w}_{\pm 2} = \left(\pm \sqrt{1 + \frac{\varepsilon^2 s}{D}}, 1, 0, 0, 0 \right), \\ \frac{\varepsilon s}{v^*}, & \quad \text{with eigenvector } \mathbf{w}_3 = (0, 0, 1, 0, 0). \end{aligned}$$

If $|s| \ll \frac{1}{\varepsilon}$, then the matrix $\widehat{\mathbf{M}}_0(\varepsilon, s)$ is not hyperbolic at $\varepsilon = 0$. However, this problem is not fundamental as it is known that there is a hyperbolic splitting in the full problem (see Sect. 3.2) and there is a spectral gap between the positive and negative eigenvalues, even if ε is close to zero. The spectral gap disappears if $s \approx -Dk^2$ (or $\varepsilon^2 s \approx -D$) and in this case the following arguments will not work. The spectral gap allows us to define a weight function which moves the spectrum away from the imaginary axis. To be specific, define

$$\widetilde{\mathbf{w}}(\eta) = e^{v\eta} \widehat{\mathbf{w}}(\eta), \quad \text{with } v = \begin{cases} \frac{1}{2} \operatorname{sgn}(s), & \text{if } |\varepsilon s| < v^*, \\ 0, & \text{if } |\varepsilon s| \geq v^*. \end{cases}$$

Then $\widetilde{\mathbf{w}}(\eta)$ satisfies the ODE

$$\partial_\eta \widetilde{\mathbf{w}} = \left([\widehat{\mathbf{M}}_0(\varepsilon, s) + v\mathbf{I}] + \varepsilon \widehat{\mathbf{M}}_1(\eta; E_\infty, \varepsilon) \right) \widetilde{\mathbf{w}}, \quad (5.12)$$

and the spectrum of $\widehat{\mathbf{M}}_0(\varepsilon, s) + v\mathbf{I}$ is bounded away from zero for all ε small (as long as $s + Dk^2$ is not small). The system

$$\partial_\eta \widetilde{\mathbf{w}} = [\widehat{\mathbf{M}}_0(\varepsilon, s) + v\mathbf{I}] \widetilde{\mathbf{w}} \quad (5.13)$$

has an exponential dichotomy with projection \mathbf{P}_0 such that the range of \mathbf{P}_0 is the span of all eigenvectors of the negative eigenvalues and the kernel of \mathbf{P}_0 is the span of all eigenvectors of the positive eigenvalues. Then \mathbf{P}_0 is the projection on the stable subspace of the linear system (5.13) and $\mathbf{I} - \mathbf{P}_0$ is the projection on the unstable subspace.

Clearly the matrix $\widehat{\mathbf{M}}_1(\eta; E_\infty, \varepsilon)$ is uniformly bounded for all η and ε small. Thus applying the Roughness Theorem 2 gives that, for small ε , there is an exponential dichotomy for the system (5.12) on \mathbb{R}^+ with projection $\mathbf{P}_\varepsilon^s(\eta)$ onto the stable subspace such that $\mathbf{P}_\varepsilon^s(\eta)$ is ε -close to \mathbf{P}_0 for all $\eta \geq 0$. And similarly, there is an exponential dichotomy for the system (5.12) on \mathbb{R}^- with projection $\mathbf{P}_\varepsilon^u(\eta)$ onto the unstable subspace such that $\mathbf{P}_\varepsilon^u(\eta)$ is ε -close to $\mathbf{I} - \mathbf{P}_0$ for all $\eta \leq 0$. Thus the range of $\mathbf{P}_\varepsilon^s(0)$ is ε -close to the range of \mathbf{P}_0 and the range of $\mathbf{P}_\varepsilon^u(0)$ is ε -close to the range of $\mathbf{I} - \mathbf{P}_0$, so the range of $\mathbf{P}_\varepsilon^s(0)$ and the range of $\mathbf{P}_\varepsilon^u(0)$ have only a trivial intersection for ε small.

As the weight function $e^{v\eta}$ has been chosen such that no eigenvalue crosses the imaginary axis, it affects only the value of the dichotomy exponentials, not their sign, nor the stable and unstable manifolds. Hence the stable and unstable manifolds of (5.11) have a trivial intersection only. And the same holds for the stable and unstable manifolds of (3.3), as the only difference between the systems (5.11) and (3.3)

is a scaling. So it can be concluded that the linear system (3.3) does not have any bounded solutions for ε small (k large) and s not close to $-Dk^2$.

If s is close to $-Dk^2$, i.e., $s = -\frac{D}{\varepsilon^2}(1 + o(1))$, then the matrix $\hat{A}_0(\varepsilon, s)$ has a positive and negative eigenvalue of order $o(1)$ and the spectral gap will disappear in the limit $\varepsilon \rightarrow 0$. So the roughness theorem cannot be applied anymore and no conclusion about bounded solutions can be drawn.

The arguments above show that only if $s = -Dk^2(1 + o(1))$ is there a possibility for bounded solutions to exist. As the dispersion curve indicates a bounded solution of the ODE (3.3), this implies that for ε near zero, hence k large, the dispersion curve is

$$s(k) = -Dk^2(1 + o(1)), \quad k \rightarrow \infty. \quad (5.14)$$

So far we have used that $s + Dk^2 > 0$. If one considers the linear system (3.3), an edge of the continuous spectrum is given by the curve $s = -Dk^2 + f(|E_\infty|)$. However, one should include the decay condition (2.11), i.e., the scaling (3.7). For any $0 < \beta < \Lambda^*$ the edge becomes $s = -Dk^2 + \beta$. By taking the limit for $\beta \rightarrow 0$, we see that the curve $s = -Dk^2$ is an edge of the continuous spectrum. Thus with the decay condition, either the dispersion curve satisfies $s(k) \geq -Dk^2$ or it ends at the continuous spectrum.

6 Physically Guided Fits to the Numerical Dispersion Relations

In Sect. 3 we derived dispersion relations for a number of fields E_∞ and diffusion constants D by numerically solving an eigenvalue problem. We confirmed these calculations through numerical solutions of the initial value problem in Sect. 4, and we derived analytical asymptotic limits to these dispersion relations in Sect. 5. This sets the stage for comparing the numerical results to the analytical asymptotic limits and for deriving physically guided empirical fits to the numerical dispersion curves where the analytical asymptotic limits are not applicable. The small k -data derived in Sect. 3.4 and the analysis of Sect. 5.1 are shown to be consistent in Sect. 6.1. After showing in Sect. 6.2 that a simple cross-over formula joining the asymptotic behavior of the small wave numbers with the asymptotic behavior of the large wave numbers is not satisfactory, we give a data collapse, empirical fits and arguments on relevant scales in Sect. 6.3.

6.1 Testing the Small k Asymptotic Limits

First the asymptotic relation (5.2) for small k is tested on the numerical results. Beyond the results visible in the plots of Fig. 2, the numerical dispersion relation for $E_\infty = -1$ and $D = 0.1$ was evaluated carefully for small values of k , and the result is shown in a double logarithmic plot in Fig. 5 that zooms in on the small k behavior. Also plotted is the analytical asymptotic limit (5.2). The comparison between numerical data and analytical asymptotic limit is convincing in the range of small k .

For the other values of E_∞ and D presented in Fig. 2, the dispersion relation $s(k)$ again is fitted very well by the asymptotic limit (5.2) for small values of k . This will be illustrated in Fig. 6.

Fig. 5 A log–log plot of the dispersion curve $s(k)$ for $E = -1$ and $D = 0.1$ illustrates the behavior for small k . *Solid line with squares*: Data from the numerical evaluation of the eigenvalue problem. *Dashed line*: Analytical asymptotic limit $\log s = \log c^* + \log k$ according to (5.2)





Fig. 6 Labeled curves: The numerical dispersion curves from Fig. 2 plotted as $\mathcal{S} = Ds/c^{*2}$ over $\kappa = Dk/c^*$. *Dotted line on the left*: the parabola $\kappa - \kappa^2$ that would be predicted by (6.1) as far as it fits into the plotted region



6.2 Testing Both Asymptotic Limits


It is quite suggestive to join the small k asymptotic limit (5.2) with the large k asymptotic limit (5.14) into one cross-over formula

$$s(k) = c^*k - Dk^2, \quad c^*(E_\infty, D) = E \left. \frac{dv^*}{dE} \right|_{E_\infty} = |E_\infty| \left(1 + f'(|E_\infty|) \sqrt{\frac{D}{f(|E_\infty|)}} \right). \quad (6.1)$$

A formula similar to $s(k) = c^*k - Dk^2$ was suggested in Arrayás et al. (2005), but with a different prefactor instead of c^* . However, we now confirm once more the correctness of the prefactor c^* , and show that the large k asymptotic limit is not yet applicable in the range of positive growth rates $s(k)$.

If (6.1) holds, then the dispersion relation for the rescaled variable $\mathcal{S} = Ds/c^{*2}$ as a function of the rescaled wave number $\kappa = Dk/c^*$ becomes $\mathcal{S} = \kappa - \kappa^2$. Therefore the formula (6.1) can easily be tested on the numerical data from Fig. 2 by plotting them in rescaled variables \mathcal{S} and κ with appropriate values for D and $c^*(E_\infty, D)$

Fig. 7 The numerical dispersion curves from Fig. 2 plotted as s/s_{\max} over k/k_0 ; here $s_{\max} = \max_k s(k)$ and $k_0 > 0$ with $s(k_0) = 0$ are determined from the respective curve



for each curve. The result is shown in Fig. 6, together with the parabola $\kappa - \kappa^2$. The plot illustrates that the asymptotic limit (5.2) indeed is a very good fit to all data for small k .

For larger k , the curves differ quantitatively. In particular, \mathcal{S} vanishes for κ between 0.014 and 0.035 for the numerical dispersion curves while the formula (6.1) predicts this to happen for $\kappa = 1$. Also the maximum of the dispersion curve \mathcal{S}_{\max} is never higher than 0.0027 for the numerical data while formula (6.1) predicts 0.25. Of course, this is not in contradiction with the analytical results in Sect. 5.3 for large k . Rather it says that the positive part of the dispersion curve lies completely in the range of small k , where the asymptotic limit for large k is not applicable. We conclude that cross-over formula (6.1) is not an appropriate fit for the numerically derived dispersion relations.

6.3 Data Collapse, Relevant Length Scales, Empirical Fits and Conjectures

We finish this section with a data collapse and arguments on relevant scales that guide empirical fits.

6.3.1 Data Collapse

First, we investigate whether the numerical data for different E_∞ and D can be collapsed onto one curve. This is done by determining the maximum of the dispersion curve s_{\max} and the wave number k_0 where the growth rate vanishes, $s(k_0) = 0$, from the numerical data for each pair (E_∞, D) . In Fig. 7 all curves are plotted as s/s_{\max} and k/k_0 with their respective s_{\max} and k_0 . The plot shows that the curve shapes are very similar, but they do not coincide completely. For example, there seems to be a small drift in the position of the maximum.

6.3.2 Relevant Length Scales and the $D = 0$ Case

In a second step, we investigate which physical or mathematical mechanisms can suppress the growth rate $s(k)$ for much smaller values of k than suggested

by the large k asymptotic limit (5.14). In a first overview, there are three length scales in the problem. The transversal perturbation is characterized by its wavelength $2\pi/k$. In the longitudinal direction, the front is characterized by two length scales, the electric screening length ℓ_α and the diffusion length ℓ_D , cf. Fig. 1,

$$\ell_\alpha = \frac{1}{\alpha(E_\infty)} \quad \text{and} \quad \ell_D = \frac{1}{\Lambda^*} = \sqrt{\frac{D}{f(E_\infty)}}. \quad (6.2)$$

For vanishing diffusion D , the diffusion length vanishes and the screening length ℓ_α has to be compared to the wavelength of the perturbation; in Arrayás and Ebert (2004) it was shown that it determines the cross-over from the small to the large k asymptotic limit of the dispersion relation:

$$s(k) = \begin{cases} |E_\infty||k|, & \text{for } |k| \ll k_\alpha, \\ |E_\infty|k_\alpha, & \text{for } |k| \gg k_\alpha; \end{cases} \quad \text{for } D = 0, \text{ where } k_\alpha = \frac{1}{2\ell_\alpha}. \quad (6.3)$$

The actual curve for $D = 0$ and $E_\infty = -1$ is given in Fig. 2b, where we remark that the form

$$s(k) = \frac{|E_\infty|k}{[1 + (\frac{k}{k_\alpha})^p]^{1/p}} \quad (6.4)$$

for positive real p reproduces the asymptotic limits (6.3), but does not fit the full numerical curve for $E_\infty = -1$ satisfactorily for any power p . The functional form of (6.4) will serve below as an inspiration for our empirical fits for $D > 0$.

Searching for why (6.4) does not properly fit the data, one realizes that there are actually two different definitions of the screening length possible:


$$\frac{1}{\ell_\alpha^+} = \alpha(E_\infty) \quad \text{and} \quad \frac{1}{\ell_\alpha^-} = \lambda^- = \int_0^{|E_\infty|} \frac{\alpha(x) dx}{|E_\infty|} + \mathcal{O}(\sqrt{D}), \quad (6.5)$$

with λ^- from (2.17). The dimensions of both quantities are the same, and they approach each other if α is constant in a large part of the integration interval $[0, |E_\infty|]$; this is the case with the Townsend approximation $\alpha(x) = e^{-1/x}$ for $|E_\infty| \gg 1$. Otherwise, ℓ_α^+ characterizes the slopes of the fields near the discontinuity of σ (Arrayás and Ebert 2004), while ℓ_α^- characterizes the decay (2.14) of the fields far behind the front for $\xi \rightarrow -\infty$. The analysis in Arrayás and Ebert (2004) shows that linear perturbations with wave numbers $k \gg 1$ couple to the inner local structure of the front and are dominated by ℓ_α^+ , while smaller k could couple to the larger spatial structure characterized by ℓ_α^- . In the following, this conjecture will be tested on the numerical data and the results show a need for future analysis.


6.3.3 Scales and Fits for $D > 0$

When diffusion is included, the diffusion length ℓ_D emerges as another length scale in the front. As illustrated in Fig. 1, instead of the discontinuous electron density in the front for $D = 0$, a diffusive layer of width $\ell_D = 1/\Lambda^*$ (2.13) builds up in

Fig. 8 The data of Fig. 2 plotted as $s = s/(\Lambda^* c^*)$ over $\mathfrak{K} = k/\Lambda^*$. The lines are given by the empirical formula (6.8)



(a) Using $a = 3/\alpha(|E_\infty|)$ in formula (6.8)



(b) Using $a = 3c^*/f(|E_\infty|)$ in formula (6.8)

the leading edge. While D increases, the dispersion relation decreases as shown in Fig. 2b. As the diffusive layer is the main new feature of the front for $D > 0$, it is plausible that the different behavior of s is created within this boundary layer. The physical mechanism is that diffusion can smear perturbations of short wavelength out, hence suppressing their growth. This process mainly takes place in the diffusive layer because gradients are largest in this region. This idea inspired an attempt in Arrayás et al. (2005) to calculate $s(k)$ by local analysis within the diffusion layer. In principle, such an approach combined with proper matched asymptotic expansions could work. However, the calculation in Arrayás et al. (2005) was intrinsically inconsistent (Ebert and Derks 2008), disagrees with our asymptotic limit for small k and therefore fits the numerical results even worse than formula (6.1), cf. Fig. 6.

We tested whether the diffusion length $\ell_D = 1/\Lambda^*$ plays a role in the dispersion relation by plotting the numerical data from Fig. 2; this time for the rescaled variables $s = s/(c^* \Lambda^*)$ over $\mathfrak{K} = k/\Lambda^*$. The result is shown in Fig. 8. It shows that the

numerical dispersion curves are well approximated by

$$s(k) \approx c^*k + \mathcal{O}(k^2), \quad \text{for } k \rightarrow 0, \quad (6.6)$$

$$k_0 \approx \Lambda^*/4, \quad \text{where } s(k_0) = 0. \quad (6.7)$$

The numerical evidence from Fig. 8 summarized in (6.7) together with the physical explanation above suggest the following conjecture.

Conjecture 1 *The largest unstable wave number of the Laplacian instability is proportional to the inverse diffusion length.*

We remark that the data give $k_0 \approx 1/(4\ell_D)$, while the cross-over formula (6.1) would suggest that $k_0 \approx \ell_\alpha/\ell_D^2$, highlighting again its inadequacy for intermediate k values. Figure 8 also shows that the value of the wave number for which the maximum of the dispersion curve is attained is in the range of $k_{\max} = 0.22k_0$ to $0.30k_0$.

The data in Fig. 8 suggest an empirical formula of the form (for $s \geq 0$)

$$s(k) = \frac{c^*k}{1+ak} \left(1 - \frac{4k}{\Lambda^*}\right) \quad \text{or} \quad \mathfrak{s} = \frac{\mathfrak{K}}{1+a\Lambda^*\mathfrak{K}} (1-4\mathfrak{K}), \quad (6.8)$$

where the parameter a will depend on the external parameters D and E_∞ . The factor c^*k creates the correct asymptotic limit (6.6) for $k \ll 1$. The factor $(1 - 4k/\Lambda^*)$ creates the nontrivial zero of the dispersion relation at k_0 (6.7). The form of the numerator is inspired by (6.4), and the proper asymptotic limit (6.3) for large k and $D = 0$ would be reached for $a = 2\ell_\alpha^+ + \mathcal{O}(\sqrt{D})$. Obviously, the empirical formula (6.8) is not valid in the asymptotic range $k \gg 1$ where $s < 0$ and where the asymptotic behavior is given by $s \approx -Dk^2$.

The functional form of formula (6.8) is supported by the following observation. If one calculates the maximum $(\mathfrak{K}_{\max}, \mathfrak{s}_{\max})$ of (6.8), it follows that $\mathfrak{K}_{\max}^2/\mathfrak{s}_{\max} = 1/4$, independently of the value of a . (The number $1/4$ directly stems from the factor 4 in $(1 - 4\mathfrak{K})$.) This relation indeed fits the numerical curves quite well, therefore the factor 4 is supported twice independently. Relevant numerical data for this and other fits is collected in Table 1.

The value for a is less obvious. The empirical formula (6.8) gives the following relation between $a\Lambda^*$ and the maximum of curve

$$\frac{1 - 8\mathfrak{K}_{\max}}{4\mathfrak{K}_{\max}^2} = a\Lambda^* = \frac{1 - 4\sqrt{\mathfrak{s}_{\max}}}{\mathfrak{s}_{\max}}.$$

The empirical values for those quotients are given in Table 1.

The limit of $D = 0$ and $k \gg 1$ suggests $a = 2\ell_\alpha^+ + \mathcal{O}(\sqrt{D})$, but the fit is unconvincing (cf. also the earlier discussion for $D = 0$). However, we found that $a = 3\ell_\alpha^+$ fits the data reasonably well. Formula (6.8) with this value of a together with the numerical data are visualized in Fig. 8a. The fit is quite good for the lower two curves, but the upper two display some discrepancies. The main problem is the value of \mathfrak{s}_{\max} with a relative error between 2% and 24%, while the position of \mathfrak{K}_{\max} has a relative error as low as 0.5% to 7%. As $f(|E_\infty|)/c^* = \alpha(|E_\infty|) + \mathcal{O}(\sqrt{D})$, another possible

Table 1 Upper block: characteristic numerical data of the dispersion relations in Fig. 2 with errors. Middle block: characteristic scales of the planar front according to analysis. Lower block: relevant ratios of numerical and analytical scales as used for the derivation of the empirical formula

(E_∞, D)	$(-1, 0.01)$	$(-1, 0.1)$	$(-2, 0.1)$	$(-5, 0.1)$	$(-10, 0.1)$
s_{\max}	0.080(1)	0.05190(2)	0.1695(15)	0.647(1)	1.6305(15)
k_{\max}	0.35(4)	0.144(1)	0.25(4)	0.45(4)	0.60(4)
k_0	1.575(15)	0.4825(5)	0.875(25)	1.595(1)	2.397(1)
k_0/k_{\max}	4.56(56)	3.35(3)	3.60(65)	3.57(33)	4.01(27)
v^*	1.12	1.38	2.70	6.28	11.9
c^*	1.12	1.38	2.52	5.77	11.0
$\Lambda^* = \sqrt{f(E_\infty)/D}$	6.07	1.92	3.48	6.40	9.51
$\alpha(E_\infty) = 1/\ell_\alpha^+$	0.37	0.37	0.61	0.82	0.90
σ^-	0.148	0.144	0.638	2.832	7.169
$\lambda^- = 1/\ell_\alpha^-$	0.13	0.10	0.23	0.45	0.60
$3\Lambda^*/\alpha(E_\infty)$	49.5	15.6	17.2	23.4	31.5
$3\Lambda^*c^*/f(E_\infty)$	55.5	21.6	21.7	27.0	34.8
$\mathfrak{R}_0 = k_0/\Lambda^*$	0.260(3)	0.252(1)	0.251(5)	0.249(1)	0.252(1)
$\mathfrak{s}_{\max} = s_{\max}/(c^*\Lambda^*)$	0.0118(2)	0.0196(1)	0.0193(2)	0.0175(1)	0.0155(1)
k_{\max}/k_α	1.9(2)	0.78(1)	0.8(2)	1.1(1)	1.3(1)
$\mathfrak{R}_{\max}^2/\mathfrak{s}_{\max}$	0.28(7)	0.288(4)	0.27(9)	0.28(5)	0.26(4)
$(1 - 8\mathfrak{R}_{\max})/(4\mathfrak{R}_{\max}^2)$	40(6)	17.7(1)	21(2)	22(2)	31(2)
$(1 - 4\sqrt{\mathfrak{s}_{\max}})/\mathfrak{s}_{\max}$	48.1(4)	22.5(1)	23.0(1)	26.8(1)	32.3(1)

fit is $a = 3c^*/f(|E_\infty|)$; it is displayed in Fig. 8b. The fit is quite good for the upper two curves, but now the fit has some discrepancies for the lower two curves. And the value of \mathfrak{R}_{\max} has a larger error between 2% and 10% while the position of \mathfrak{s}_{\max} has a much smaller error of only 1% to 10%. Obviously, these observations require further analytical investigation. Note finally the striking relation between $\lambda^- = 1/\ell_\alpha^-$ and the value of k_{\max} for larger values of the electric field in Table 1. As a basis for future work, all characteristic numerical data are collected in this table.

7 Conclusion and Outlook

In this paper, we found dispersion curves for negative streamer ionization fronts by numerically solving an eigenvalue problem. We verified this prediction on the numerical solution of an initial value problem, we derived analytical expressions for the asymptotics of the curve for large and small wavelengths, and we presented a physically motivated fit formula to the numerical curves for intermediate wavelengths. The investigation is of interest for two reasons: because pulled fronts like these are mathematically challenging to investigate, and because explicit predictions on the linear

stability of ionization fronts help to interpret numerical and experimental observations of propagating and branching streamer discharges.

The ionization front is a pulled front, i.e., the front is part of a family of traveling waves that propagate into a temporally unstable steady state. For the dynamics with one spatial variable, most traveling waves in this family are attractors only for waves with exactly the same asymptotic decay profile. The exception is the pulled front, which has the steepest decay of all waves in the family and is an attractor for waves with a sufficiently fast decay (therefore excluding the slower decay rates for the other traveling waves in the family). The instability of the state ahead of the front and the related spatial decay condition imply that only a submanifold of the stable manifold in the transverse instability problem is relevant for the transverse instability analysis. This submanifold is identified by introducing a weighted solution space that excludes solutions with a decay rate that is too slow. We integrated the relevant stable submanifold and unstable manifold numerically with a dynamical systems method to calculate the dispersion curve. This method of finding the dispersion curve does not use any details of the streamer model, except that it has a pulled front. The definition of a submanifold of the stable manifold and the subsequent numerical integration of this stable submanifold and the unstable manifold are ideas that can be applied to pulled fronts in other systems, too.

It is interesting to see that the band of unstable wave numbers seems to be limited by a multiple of the decay rate Λ^* that characterizes the leading edge of the pulled ionization front; though the evidence up to now is only numerical. As such behavior is physically reasonable, the next step would be to derive it analytically, e.g., by a local analysis in the diffusive layer and matched asymptotic expansions. Such an expansion could be based on the limiting case where the diffusion length $\ell_D = 1/\Lambda^*$ is much smaller than the screening length ℓ_α .

The calculated dispersion curves also contribute to understanding the stability of actual streamers. Two- and three-dimensional time-dependent simulations (Arrayás et al. 2002; Rocco et al. 2002; Montijn et al. 2006a, 2006b; Luque et al. 2007) of the streamer model introduced in Sect. 2 show them to become unstable and branch. Can the unstable wavelengths of this branching be related to the unstable band of wavelengths of the present calculation? Furthermore, if the inner front structure is approximated by a moving boundary (Meulenbroek et al. 2005; Ebert et al. 2007), how is the calculated dispersion relation of transversal perturbations to be taken into account? It will also be interesting to see whether the dispersion relation calculated for the present fluid model is also applicable to the corresponding particle model (Li et al. 2007).

Finally, we mention that the extension of the streamer model with photo-ionization as an additional reaction term (Luque et al. 2007) in composed gases like air requires an extension of the present analysis as nonlocal interaction terms play a role.

Appendix: Matrices in Exterior Algebra Spaces

In this appendix, we give explicit expressions for the matrices $\mathbf{M}^{(l)}$ acting on the exterior algebra space $\bigwedge^l(\mathbb{C}^5)$ for $l = 2, 3$. Let $\mathbf{e}_1, \dots, \mathbf{e}_5$ be the standard basis for \mathbb{C}^5 .

Then an induced basis on $\bigwedge^2(\mathbb{C}^5)$ is given by

$$\begin{aligned} \mathbf{a}_1 &= \mathbf{e}_1 \wedge \mathbf{e}_2, & \mathbf{a}_2 &= \mathbf{e}_1 \wedge \mathbf{e}_3, & \mathbf{a}_3 &= \mathbf{e}_1 \wedge \mathbf{e}_4, & \mathbf{a}_4 &= \mathbf{e}_1 \wedge \mathbf{e}_5, & \mathbf{a}_5 &= \mathbf{e}_2 \wedge \mathbf{e}_3, \\ \mathbf{a}_6 &= \mathbf{e}_2 \wedge \mathbf{e}_4, & \mathbf{a}_7 &= \mathbf{e}_2 \wedge \mathbf{e}_5, & \mathbf{a}_8 &= \mathbf{e}_3 \wedge \mathbf{e}_4, & \mathbf{a}_9 &= \mathbf{e}_3 \wedge \mathbf{e}_5, & \mathbf{a}_{10} &= \mathbf{e}_4 \wedge \mathbf{e}_5. \end{aligned}$$

The matrix $\mathbf{M}^{(2)} : \bigwedge^2(\mathbb{C}^5) \rightarrow \bigwedge^2(\mathbb{C}^5)$ can be associated with a complex 10×10 matrix with entries such that

$$\mathbf{M}^{(2)} \mathbf{a}_i = \sum_{j=1}^{10} \mathbf{M}_{ij}^{(2)} \mathbf{a}_j, \quad i, j = 1, \dots, 10, \quad (8.1)$$

where, for any decomposable $\mathbf{x} = \mathbf{x}_1 \wedge \mathbf{x}_2 \in \bigwedge^2(\mathbb{C}^5)$, $\mathbf{M}^{(2)} \mathbf{x} := \mathbf{M} \mathbf{x}_1 \wedge \mathbf{x}_2 + \mathbf{x}_1 \wedge \mathbf{M} \mathbf{x}_2$. Let \mathbf{M} be an arbitrary 5×5 matrix with complex entries,

$$\mathbf{M} = \begin{pmatrix} m_{11} & m_{12} & m_{13} & m_{14} & m_{15} \\ m_{21} & m_{22} & m_{23} & m_{24} & m_{25} \\ m_{31} & m_{32} & m_{33} & m_{34} & m_{35} \\ m_{41} & m_{42} & m_{43} & m_{44} & m_{45} \\ m_{51} & m_{52} & m_{53} & m_{54} & m_{55} \end{pmatrix}, \quad (8.2)$$

then $\mathbf{M}^{(2)}$ takes the explicit form

$$\mathbf{M}^{(2)} = \begin{bmatrix} d_{12} & m_{23} & m_{24} & m_{25} & -m_{13} & -m_{14} & -m_{15} & 0 & 0 & 0 \\ m_{32} & d_{13} & m_{34} & m_{35} & m_{12} & 0 & 0 & -m_{14} & -m_{15} & 0 \\ m_{42} & m_{43} & d_{14} & m_{45} & 0 & m_{12} & 0 & m_{13} & 0 & -m_{15} \\ m_{52} & m_{53} & m_{54} & d_{15} & 0 & 0 & m_{12} & 0 & m_{13} & m_{14} \\ -m_{31} & m_{21} & 0 & 0 & d_{23} & m_{34} & m_{35} & -m_{24} & -m_{25} & 0 \\ -m_{41} & 0 & m_{21} & 0 & m_{43} & d_{24} & m_{45} & m_{23} & 0 & -m_{25} \\ -m_{51} & 0 & 0 & m_{21} & m_{53} & m_{54} & d_{25} & 0 & m_{23} & m_{24} \\ 0 & -m_{41} & m_{31} & 0 & -m_{42} & m_{32} & 0 & d_{34} & m_{45} & -m_{35} \\ 0 & -m_{51} & 0 & m_{31} & -m_{52} & 0 & m_{32} & m_{54} & d_{35} & m_{34} \\ 0 & 0 & -m_{51} & m_{41} & 0 & -m_{52} & m_{42} & -m_{53} & m_{43} & d_{45} \end{bmatrix},$$

where $d_{ij} = m_{ii} + m_{jj}$.

In a similar way, the matrix $\mathbf{M}^{(3)} : \bigwedge^3(\mathbb{C}^5) \rightarrow \bigwedge^3(\mathbb{C}^5)$ can be associated with a complex 10×10 matrix. First we define an induced basis on $\bigwedge^3(\mathbb{C}^5)$ by

$$\begin{aligned} \mathbf{b}_1 &= \mathbf{e}_1 \wedge \mathbf{e}_2 \wedge \mathbf{e}_3, & \mathbf{b}_2 &= \mathbf{e}_1 \wedge \mathbf{e}_2 \wedge \mathbf{e}_4, & \mathbf{b}_3 &= \mathbf{e}_1 \wedge \mathbf{e}_2 \wedge \mathbf{e}_5, & \mathbf{b}_4 &= \mathbf{e}_1 \wedge \mathbf{e}_3 \wedge \mathbf{e}_4, \\ \mathbf{b}_5 &= \mathbf{e}_1 \wedge \mathbf{e}_3 \wedge \mathbf{e}_5, & \mathbf{b}_6 &= \mathbf{e}_1 \wedge \mathbf{e}_4 \wedge \mathbf{e}_5, & \mathbf{b}_7 &= \mathbf{e}_2 \wedge \mathbf{e}_3 \wedge \mathbf{e}_4, & \mathbf{b}_8 &= \mathbf{e}_2 \wedge \mathbf{e}_3 \wedge \mathbf{e}_5, \\ \mathbf{b}_9 &= \mathbf{e}_2 \wedge \mathbf{e}_4 \wedge \mathbf{e}_5, & \mathbf{b}_{10} &= \mathbf{e}_3 \wedge \mathbf{e}_4 \wedge \mathbf{e}_5. \end{aligned}$$

The matrix for $\mathbf{M}^{(3)} \in \mathbb{C}^{10 \times 10}$ has entries such that

$$\mathbf{M}^{(3)} \mathbf{b}_i = \sum_{j=1}^{10} \mathbf{M}_{ij}^{(3)} \mathbf{b}_j, \quad i, j = 1, \dots, 10, \quad (8.3)$$

where, for any decomposable $\mathbf{x} = \mathbf{x}_1 \wedge \mathbf{x}_2 \wedge \mathbf{x}_3 \in \bigwedge^3(\mathbb{C}^5)$, $\mathbf{M}^{(3)}\mathbf{x} := \mathbf{M}\mathbf{x}_1 \wedge \mathbf{x}_2 \wedge \mathbf{x}_3 + \mathbf{x}_1 \wedge \mathbf{M}\mathbf{x}_2 \wedge \mathbf{x}_3 + \mathbf{x}_1 \wedge \mathbf{x}_2 \wedge \mathbf{M}\mathbf{x}_3$. If M is given by (8.2), then $\mathbf{M}^{(3)}$ takes the explicit form

$$\mathbf{M}^{(3)} = \begin{bmatrix} d_{123} & m_{43} & m_{53} & -m_{42} & -m_{52} & 0 & m_{41} & m_{51} & 0 & 0 \\ m_{34} & d_{124} & m_{54} & m_{32} & 0 & -m_{52} & -m_{31} & 0 & m_{51} & 0 \\ m_{35} & m_{45} & d_{125} & 0 & m_{32} & m_{42} & 0 & -m_{31} & -m_{41} & 0 \\ -m_{24} & m_{23} & 0 & d_{134} & m_{54} & -m_{53} & m_{21} & 0 & 0 & m_{51} \\ -m_{25} & 0 & m_{23} & m_{45} & d_{135} & m_{43} & 0 & m_{21} & 0 & -m_{41} \\ 0 & -m_{25} & m_{24} & -m_{35} & m_{34} & d_{145} & 0 & 0 & m_{21} & m_{31} \\ m_{14} & -m_{13} & 0 & m_{12} & 0 & 0 & d_{234} & m_{54} & -m_{53} & m_{52} \\ m_{15} & 0 & -m_{13} & 0 & m_{12} & 0 & m_{45} & d_{235} & m_{43} & -m_{42} \\ 0 & m_{15} & -m_{14} & 0 & 0 & m_{12} & -m_{35} & m_{34} & d_{245} & m_{32} \\ 0 & 0 & 0 & m_{15} & -m_{14} & m_{13} & m_{25} & -m_{24} & m_{23} & d_{345} \end{bmatrix},$$

where $d_{ijk} = m_{ii} + m_{jj} + m_{kk}$.

Acknowledgements We thank Björn Sandstede for helpful discussions on the Roughness Theorem. We thank Willem Hundsdorfer and René Reimer for advice and help with the simulations in Sect. 4.

References

- Alexander, J., Gardner, R., Jones, C.K.R.T.: A topological invariant arising in the stability analysis of traveling waves. *J. Reine Angew. Math.* **410**, 167–212 (1990)
- Allen, L., Bridges, T.J.: Numerical exterior algebra and the compound matrix method. *Numer. Math.* **92**, 197–232 (2002)
- Arrayás, M., Ebert, U.: Stability of negative ionization fronts: Regularization by electric screening? *Phys. Rev. E* **69**, 036214 (2004), 10 p.
- Arrayás, M., Ebert, U., Hundsdorfer, W.: Spontaneous branching of anode-directed streamers between planar electrodes. *Phys. Rev. Lett.* **88**, 174502 (2002), 4 p.
- Arrayás, M., Fontelos, M.A., Trueba, J.L.: Mechanism of branching in negative ionization fronts. *Phys. Rev. Lett.* **95**, 165001 (2005), 4 p.
- Back, A., Guckenheimer, J., Myers, M.R., Wicklin, F.J., Worfolk, P.A.: DsTool: Computer assisted exploration of dynamical systems. *Not. Amer. Math. Soc.* **39**, 303–309 (1992)
- Bawagan, A.D.O.: A stochastic model of gaseous dielectric breakdown. *Chem. Phys. Lett.* **281**, 325–331 (1997)
- Blyuss, K.B., Bridges, T.J., Derks, G.: Transverse instability and its long-term development for solitary waves of the (2+1)-dimensional Boussinesq equation. *Phys. Rev. E* **67**, 056626 (2003), 9 p.
- Brau, F., Luque, A., Meulenbroek, B., Ebert, U., Schäfer, L.: Construction and test of a moving boundary model for negative streamer discharges. *Phys. Rev. E* **77**, 026219 (2008)
- Bridges, T., Derks, G., Gottwald, G.A.: Stability and instability of solitary waves of the fifth-order KdV equation: a numerical framework. *Physica D* **172**, 190–216 (2003)
- Brin, L.Q.: Numerical testing of the stability of viscous shock waves. *Math. Comput.* **70**, 1071–1088 (2001)
- Brin, L.Q., Zumbrun, K.: Analytically varying eigenvectors and the stability of viscous shock waves. In: *Seventh Workshop on Partial Differential Equations, Part I*, Rio de Janeiro, 2001. *Mat. Contemp.*, vol. 22, pp. 19–32. Instituto de Matematica Pura e Aplicada, Rio de Janeiro (2002)
- Coppel, W.: *Dichotomies in Stability Theory*, Lecture Notes in Mathematics. Springer, Berlin (1978)

- Derks, G., Gottwald, G.A.: A robust numerical method to study oscillatory instability of gap solitary waves. *SIAM J. Appl. Dyn. Sys.* **4**, 140–158 (2005)
- Dhali, S.K., Williams, P.F.: Numerical simulation of streamer propagation in nitrogen at atmospheric pressure. *Phys. Rev. A* **31**, 1219–1221 (1985)
- Dhali, S.K., Williams, P.F.: Two-dimensional studies of streamers in gases. *J. Appl. Phys.* **62**, 4696–4707 (1987)
- Ebert, U., Arrayás, M.: Pattern formation in electric discharges. In: Reguera, D., et al. (eds.) *Coherent Structures in Complex Systems. Lecture Notes in Physics*, vol. 567, pp. 270–282. Springer, Berlin (2001)
- Ebert, U., Derks, G.: Comment on Arrayás et al. (2005). *Phys. Rev. Lett.* (2008, submitted), 1 p.
- Ebert, U., Meulenbroek, B., Schäfer, L.: Rigorous stability results for a Laplacian moving boundary problem with kinetic undercooling. *SIAM J. Appl. Math.* **69**, 292–310 (2007)
- Ebert, U., Montijn, C., Briels, T.M.P., Hundsdoerfer, W., Meulenbroek, B., Rocco, A., van Veldhuizen, E.M.: The multiscale nature of streamers. *Plasma Sources Sci. Technol.* **15**, S118–S129 (2006)
- Ebert, U., van Saarloos, W.: Universal algebraic relaxation of fronts propagating into an unstable state and implications for moving boundary approximations. *Phys. Rev. Lett.* **80**, 1650–1653 (1998)
- Ebert, U., van Saarloos, W.: Front propagation into unstable states: Universal algebraic convergence towards uniformly translating pulled fronts. *Physica D* **146**, 1–99 (2000a)
- Ebert, U., van Saarloos, W.: Breakdown of the standard perturbation theory and moving boundary approximation for “pulled” fronts. *Phys. Rep.* **337**, 139–156 (2000b)
- Ebert, U., van Saarloos, W., Caroli, C.: Streamer propagation as a pattern formation problem: planar fronts. *Phys. Rev. Lett.* **77**, 4178–4181 (1996)
- Ebert, U., van Saarloos, W., Caroli, C.: Propagation and structure of planar streamer fronts. *Phys. Rev. E* **55**, 1530–1594 (1997)
- Evans, J.W.: Nerve axon equations IV. The stable and unstable impulse. *Indiana Univ. Math. J.* **24**, 1169–1190 (1975)
- Humpherys, J., Zumbrun, K.: An efficient shooting algorithm for Evans function calculations in large systems. *Physica D* **220**, 116–126 (2006)
- Humpherys, J., Sandstede, B., Zumbrun, K.: Efficient computation of analytic bases in Evans function analysis of large systems. *Numer. Math.* **103**, 631–642 (2006)
- Kapitula, T.: The Evans function and generalized Melnikov integrals. *SIAM J. Math. Anal.* **30**, 273–297 (1999)
- Kapitula, T., Sandstede, B.: Eigenvalues and resonances using the Evans function. *Discrete Contin. Dyn. Syst.* **10**, 857–869 (2004)
- Kato, T.: *Perturbation Theory for Linear Operators*, 2nd edn. Springer, Heidelberg (1984)
- Li, C., Brok, W.J.M., Ebert, U., van der Mullen, J.J.A.M.: Deviations from the local field approximation in negative streamer heads. *J. Appl. Phys.* **101**, 123305 (2007), 14 p.
- Luque, A., Ebert, U., Montijn, C., Hundsdoerfer, W.: Photoionisation in negative streamers: fast computations and two propagation modes. *Appl. Phys. Lett.* **90**, 081501 (2007), 3 p.
- Marcus, M.: *Finite Dimensional Multilinear Algebra, Part II*. Dekker, New York (1975)
- Meulenbroek, B., Ebert, U., Schäfer, L.: Regularization of moving boundaries in a Laplacian field by a mixed Dirichlet–Neumann boundary condition: exact results. *Phys. Rev. Lett.* **95**, 195004 (2005), 4 p.
- Montijn, C., Ebert, U., Hundsdoerfer, W.: Numerical convergence of the branching time of negative streamers. *Phys. Rev. E* **73**, 065401 (2006a), 4 p.
- Montijn, C., Hundsdoerfer, W., Ebert, U.: An adaptive grid refinement strategy for the simulation of negative streamers. *J. Comput. Phys.* **219**, 801–835 (2006b)
- Niemeyer, L., Pietronero, L., Wiesman, H.J.: Fractal dimension of dielectric breakdown. *Phys. Rev. Lett.* **52**, 1033–1036 (1984)
- Niemeyer, L., Ullrich, L., Wiegart, N.: The mechanism of leader breakdown in electronegative gases. *IEEE Trans. Electr. Insul.* **24**, 309–324 (1989)
- Pasko, V.P., Inan, U.S., Bell, T.F.: Mesosphere–troposphere coupling due to sprites. *Geophys. Res. Lett.* **28**, 3821–3824 (2001)
- Peterhof, D., Sandstede, B., Scheel, A.: Exponential dichotomies for solitary-wave solutions of semilinear elliptic equations on infinite cylinders. *J. Differ. Equ.* **140**, 266–308 (1997)
- Rocco, A., Ebert, U., Hundsdoerfer, W.: Branching of negative streamers in free flight. *Phys. Rev. E* **66**, 035102(R) (2002), 4 p.
- Rodin, P., Ebert, U., Hundsdoerfer, W., Grekhov, I.V.: Superfast fronts of impact ionization in initially unbiased layered semiconductor structures. *J. Appl. Phys.* **92**, 1971–1980 (2002)

- Sandstede, B.: Stability of travelling waves. In: Fiedler, B. (ed.) *Handbook of Dynamical Systems II*, pp. 983–1055. North-Holland, Amsterdam (2002)
- Starikovskaia, S.M.: Plasma assisted ignition and combustion. *J. Phys. D: Appl. Phys.* **39**, R265–R299 (2006)
- Terman, D.: Stability of planar wave solutions to a combustion model. *SIAM J. Math. Anal.* **21**, 1139–1171 (1990)
- Vitello, P.A., Penetrante, B.M., Bardsley, J.N.: Simulation of negative streamer dynamics in nitrogen. *Phys. Rev. E* **49**, 5574–5598 (1994)