

On the existence of real R-matrices for virtual link invariants

Guus Regts¹ · Alexander Schrijver¹ · Bart Sevenster¹

Received: 6 March 2015 / Published online: 4 January 2017
© The Author(s) 2016

Abstract We characterize the virtual link invariants that can be described as partition function of a real-valued R-matrix, by being weakly reflection positive. Weak reflection positivity is defined in terms of joining virtual link diagrams, which is a specialization of joining virtual link diagram tangles. Basic techniques are the first fundamental theorem of invariant theory, the Hanlon–Wales theorem on the decomposition of Brauer algebras, and the Procesi–Schwarz theorem on inequalities for closed orbits.

1 Introduction

This paper is inspired by some recent results in the range of characterizing combinatorial parameters using invariant theory, in particular by Szegedy [12] and Freedman et al. [1]. We here consider the application to virtual links, which requires some new techniques from the representation theory of the symmetric group. The concepts of virtual link diagram and virtual link were introduced by Kauffman [5]; see Manturov and Ilyutko [7] and Kauffman [6] for more background.

A *virtual link diagram* is an undirected 4-regular graph G such that at each vertex v a cyclic order of the edges incident with v is specified, together with one pair of edges opposite at v that is labeled as ‘overcrossing’. The standard way of indicating this is as

$$\begin{array}{c} \diagup \quad \diagdown \\ \times \\ \diagdown \quad \diagup \end{array} \quad \text{or just} \quad \begin{array}{c} \diagup \quad \diagdown \\ \times \\ \diagdown \quad \diagup \end{array} . \quad (1)$$

To the memory of Rudolf Halin.

The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 339109.

Alexander Schrijver
a.schrijver@uva.nl

¹ University of Amsterdam, Amsterdam, The Netherlands

Vertices of a virtual link diagram are called *crossings*. Loops and multiple edges are allowed. Moreover, the ‘unknot’ is allowed, that is, the loop \bigcirc without a crossing. Let \mathcal{G} denote the collection of virtual link diagrams, two of them being the same if they are isomorphic.

In the usual way, Reidemeister moves yield an equivalence relation on virtual link diagrams. A *virtual link* is an equivalence class of virtual link diagrams. A *virtual link invariant* is a function defined on \mathcal{G} that is invariant under Reidemeister moves. (So in fact it is a function on virtual links, but the definition as given turns out to be more convenient.)

A virtual link diagram can be seen as the projection of a link in $M \times \mathbb{R}$ on M , where M is some oriented surface. Since this connection however is not stable under all Reidemeister moves (e.g., one may need to create a handle to allow a type II Reidemeister move), we will view virtual link diagrams just abstractly as given above.

In this paper, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and for any $n \in \mathbb{Z}_+$:

$$[n] := \{1, \dots, n\}. \tag{2}$$

Choose $n \in \mathbb{Z}_+$. Let the symmetric group S_2 act on $(\mathbb{R}^n)^{\otimes 4}$ so that the nonidentity element of S_2 brings $x_1 \otimes x_2 \otimes x_3 \otimes x_4$ to $x_3 \otimes x_4 \otimes x_1 \otimes x_2$. Define

$$\mathcal{R}_n := ((\mathbb{R}^n)^{\otimes 4})^{S_2}, \tag{3}$$

which is the linear space of S_2 -invariant elements of $(\mathbb{R}^n)^{\otimes 4}$. Note that \mathcal{R}_n can be identified with the collection of symmetric matrices in $(\mathbb{R}^{n \times n})^{\otimes 2}$.

Following de la Harpe and Jones [4], we call any element R of \mathcal{R}_n a *vertex model* (‘edge-coloring model’ in [12]). For any $R \in \mathcal{R}_n$, let f_R be the *partition function* of R ; that is, f_R is the function $f_R : \mathcal{G} \rightarrow \mathbb{R}$ defined by

$$f_R(G) = \sum_{\phi: E(G) \rightarrow [n]} \prod_{v \in V(G)} R_{\phi(\delta(v))}. \tag{4}$$

Here we put

$$\phi(\delta(v)) := (\phi(e_1), \phi(e_2), \phi(e_3), \phi(e_4)), \tag{5}$$

where e_1, e_2, e_3, e_4 are the edges incident with v , in clockwise order, and where e_1, e_3 form the overcrossing pair. Since R is S_2 -invariant, $R_{\phi(\delta(v))}$ is well-defined. Note that $f_R(\bigcirc) = n$.

The well-known sufficient conditions on R for f_R to be a virtual link invariant are:



$$\begin{aligned} \text{(i)} \quad & \sum_a R_{iaaj} = \delta_{ij} \quad \text{for all } i, j, \\ \text{(ii)} \quad & \sum_{a,b} R_{ijab} R_{alkb} = \delta_{ik} \delta_{jl} \quad \text{for all } i, j, k, l, \\ \text{(iii)} \quad & \sum_{a,b,c} R_{iabh} R_{jkca} R_{bclm} = \sum_{a,b,c} R_{ijbc} R_{bkla} R_{camh} \quad \text{for all } i, j, k, l, m, h, \end{aligned} \tag{6}$$

where R is expressed in the standard basis of $(\mathbb{R}^n)^{\otimes 4}$, where all indices run from 1 to n , and where δ_{ij} is the Kronecker delta. Condition (iii) is the *Yang–Baxter equation*. In the real case, the conditions (6) are also necessary conditions for f_R to be a virtual link invariant. Elements R of \mathcal{R}_n satisfying (6) are called *R-matrices*. (Often condition (i) is deleted, to obtain an invariant for ‘ribbon links’.)

In this paper, we characterize which real-valued functions f on the collection \mathcal{G} are equal to f_R for some R-matrix R . To this end, we introduce the concept of a k -join of virtual link diagrams (for any $k \in \mathbb{Z}_+$). To define it, we consider the linear space $\mathbb{R}\mathcal{G}$ of all formal \mathbb{R} -linear combinations of elements of \mathcal{G} . Any function on \mathcal{G} to a linear space can be extended

uniquely to a linear function on $\mathbb{R}\mathcal{G}$. The elements of $\mathbb{R}\mathcal{G}$ are called *quantum virtual link diagrams*.

The k -join $G \overset{k}{\vee} H$ of virtual link diagrams G and H is an element of $\mathbb{R}\mathcal{G}$. It is obtained from the disjoint union of G and H , by taking the sum over all quantum virtual link diagrams obtained as follows: choose distinct crossings u_1, \dots, u_k of G and distinct crossings v_1, \dots, v_k of H , and for each $i = 1, \dots, k$

replace  by $\frac{1}{2} (\img alt="Diagram showing crossings u_i and v_i connected by a strand." data-bbox="521 206 634 271"/> + ).$

(7)

As usual, a circle around a crossing in these pictures means that the crossing does not correspond to a crossing of the virtual link diagram, but is an artefact of the planarity of the drawing. Note that in (7), the new connections conform to the cyclic orders and the overcrossings at u_i and v_i .

The k -join can be described in terms of joining two virtual link diagram tangles (i.e., virtual link diagrams in which labeled vertices of degree 1 are allowed) by identifying equally labeled vertices (cf. Szegedy [12]). Then the k -join is obtained by ‘opening’ G and H at the crossings $u_1, \dots, u_k, v_1, \dots, v_k$ (that is, deleting these vertices topologically, thus leaving, for each deleted vertex, four open end segments). Choosing appropriate labelings at the ends and joining the tangles along equally labeled ends, yields the k -join. The k -join is therefore a more restricted operation, which will yield therefore a stronger characterization.

We call f *weakly reflection positive* if for each $k \in \mathbb{Z}_+$, the $\mathcal{G} \times \mathcal{G}$ matrix

$$(8) \quad M_{f,k} := (f(G \overset{k}{\vee} H))_{G,H \in \mathcal{G}} \tag{8}$$

is positive semidefinite. Moreover, $f : \mathcal{G} \rightarrow \mathbb{R}$ is called *multiplicative* if $f(\emptyset) = 1$ (where \emptyset is the virtual link diagram with no crossings and edges) and $f(G \sqcup H) = f(G)f(H)$ for all virtual link diagrams G, H , where \sqcup denotes disjoint union.

Theorem *Let $f : \mathcal{G} \rightarrow \mathbb{R}$. Then there exists an R-matrix R with $f = f_R$ if and only if $f(\bigcirc) \geq 0$ and f is multiplicative and weakly reflection positive and satisfies*

$$\begin{aligned}
 \text{(i)} \quad & f(\img alt="Diagram of two linked circles with a crossing." data-bbox="248 643 448 673}) + f(\bigcirc) = 2f(\img alt="Diagram of two linked circles." data-bbox="458 643 658 673}), \\
 \text{(ii)} \quad & f(\img alt="Diagram of two linked circles with two crossings." data-bbox="248 678 498 748}) + f(\bigcirc)^2 = 2f(\img alt="Diagram of two separate circles." data-bbox="708 688 818 738}), \\
 \text{(iii)} \quad & f(\img alt="Diagram of two linked circles with four crossings." data-bbox="248 753 498 844}) = f(\img alt="Diagram of two linked circles with four crossings in a different configuration." data-bbox="568 753 818 844}).
 \end{aligned}$$

(9)

Our proof of the theorem follows the line of proof layed down in [9] for ‘3-graphs’ and cyclic cubic graphs. The main addition of the present study is the application to virtual link diagrams, which requires a different combinatorial proof for the integrality of $f(\bigcirc)$. An interesting feature for virtual link diagrams is that the multiplicativity and weak reflection

positivity of f imply that $f(\bigcirc)$ is an integer but might be negative. In fact, if $f(\bigcirc)$ is negative it is even—see the lemma below. This raises the question to classify those multiplicative and weakly reflection positive virtual link invariants f with $f(\bigcirc) < 0$.

It can also be shown, with the Stone-Weierstrass theorem as in [9], that the R-matrix R in the theorem is unique, up to the natural action of the real orthogonal group $O(n)$ on R (which action leaves f_R invariant).

Multiplicative weakly reflection positive functions $f : \mathcal{G} \rightarrow \mathbb{R}$ with $f(\bigcirc) = -2k$ do exist for any $k \in \mathbb{Z}_+$. Indeed, define $f(G) = 0$ if G has at least one crossing, and $f(G) = (-2k)^t$ if G is the disjoint union of t copies of \bigcirc . Then f trivially is multiplicative, and it is weakly reflection positive, as can be derived again from the results of Hanlon and Wales [3] displayed below.

The remainder of this paper is devoted to proving the theorem.

2 The algebra homomorphism $p_n : \mathbb{R}\mathcal{G} \rightarrow \mathcal{O}(\mathcal{R}_n)$

We make some preparations to the proof of the theorem. The space $\mathbb{R}\mathcal{G}$ of formal linear combinations of elements of \mathcal{G} , is in fact an algebra, by taking the disjoint union $G \sqcup H$ of two virtual link diagrams G and H as multiplication GH . Choose $n \in \mathbb{Z}_+$ and recall that \mathcal{R}_n denotes the linear space

$$\mathcal{R}_n := ((\mathbb{R}^n)^{\otimes 4})^{S_2}. \tag{10}$$

As usual, $\mathcal{O}(\mathcal{R}_n)$ denotes the algebra of polynomials on \mathcal{R}_n . Define an algebra homomorphism $p_n : \mathbb{R}\mathcal{G} \rightarrow \mathcal{O}(\mathcal{R}_n)$ by

$$p_n(G)(R) := f_R(G) \tag{11}$$

for $G \in \mathcal{G}$ and $R \in \mathcal{R}_n$. So the element R in the theorem can be described as a common zero of the polynomials $p_n(G) - f(G)$ for all $G \in \mathcal{G}$.

We mention a connection of the k -join of virtual link diagrams to k -th derivatives of p_n , which is similar to a lemma proved in [9] for cubic cyclic graphs, and can be proved by a word for word translation of the method.

For any $q \in \mathcal{O}(\mathcal{R}_n)$, let dq be its derivative, being an element of $\mathcal{O}(\mathcal{R}_n) \otimes \mathcal{R}_n^*$. So $d^k q \in \mathcal{O}(\mathcal{R}_n) \otimes (\mathcal{R}_n^*)^{\otimes k}$. Note that the standard inner product on \mathbb{R}^n induces an inner product on $(\mathbb{R}^n)^{\otimes 4}$, hence on \mathcal{R}_n and \mathcal{R}_n^* , and therefore it induces a product $\langle \cdot, \cdot \rangle : (\mathcal{O}(\mathcal{R}_n) \otimes (\mathcal{R}_n^*)^{\otimes k}) \times (\mathcal{O}(\mathcal{R}_n) \otimes (\mathcal{R}_n^*)^{\otimes k}) \rightarrow \mathcal{O}(\mathcal{R}_n)$. Then, for all $G, H \in \mathcal{G}$ and all $k, n \in \mathbb{Z}_+$:

$$p_n(G \overset{k}{\vee} H) = \langle d^k p_n(G), d^k p_n(H) \rangle. \tag{12}$$

This connection between k -joins and k -th derivatives will be used a number of times in our proof of the theorem.

As in [12] (cf. [2, 11]), the first fundamental theorem of invariant theory for the real orthogonal group $O(n)$ implies

$$p_n(\mathbb{R}\mathcal{G}) = \mathcal{O}(\mathcal{R}_n)^{O(n)}, \tag{13}$$

the latter denoting the space of $O(n)$ -invariant elements of $\mathcal{O}(\mathcal{R}_n)$.

3 The value of f on \bigcirc

The following lemma on $f(\bigcirc)$ carries the most combinatorial part of the proof. It is based on basic results of Hanlon and Wales [3] on the representation theory of the symmetric group (cf. Sagan [10]).

Lemma *If $f : \mathcal{G} \rightarrow \mathbb{R}$ is multiplicative and weakly reflection positive, then $f(\bigcirc)$ belongs to $\{\dots, -6, -4, -2, 0, 1, 2, 3, \dots\}$.*

Proof I. We first describe some tools, using results of [3]. Consider any $k \in \mathbb{Z}_+$. For any matching M on $[8k]$ and any $\pi \in S_{8k}$, let $\pi \cdot M$ be the matching $\{\pi(e) \mid e \in M\}$. Define \mathcal{M} to be the set of perfect matchings on $[8k]$. So the group S_{8k} acts on \mathcal{M} , which induces an action of S_{8k} on $\mathbb{R}^{\mathcal{M}}$.

To each $M \in \mathcal{M}$ we can associate a virtual link diagram G_M on $[2k]$ by identifying, for each $j \in [2k]$, the vertices $4j - 3, 4j - 2, 4j - 1, 4j$ of M to one crossing called j as in

$$\begin{array}{ccc}
 \begin{array}{c} \downarrow \\ \downarrow \\ \downarrow \\ \downarrow \\ \hline 4j-3 \quad 4j-2 \quad 4j-1 \quad 4j \end{array} & \longrightarrow & \begin{array}{c} \cup \\ \cup \\ \cup \\ \cup \\ \hline j \end{array}
 \end{array}
 \tag{14}$$

To describe $G_M \overset{2k}{\vee} G_N$ for $M, N \in \mathcal{M}$, we define the following subgroups of S_{8k} . For $j \in [2k]$, let B_j be the group consisting of the identity id and of $(4j - 3, 4j - 1)(4j - 2, 4j)$. Define $B := B_1 B_2 \dots B_{2k}$. Let D be the group of permutations $d \in S_{8k}$ for which there exists $\pi \in S_{2k}$ such that $d(4j - i) = 4\pi(j) - i$ for each $j = 1, \dots, 2k$ and $i = 0, \dots, 3$. Set $Q := BD$, which is a group.

For $M, N \in \mathcal{M}$, let $c(M, N)$ denote the number of connected components of the graph $([8k], M \cup N)$. Then, by definition of the operation $\overset{2k}{\vee}$, we have

$$G_M \overset{2k}{\vee} G_N = 2^{-2k} (2k)! \sum_{s \in Q} \bigcirc^{c(M, s \cdot N)}.
 \tag{15}$$

For $\pi \in S_{8k}$, let P_π be the $\mathcal{M} \times \mathcal{M}$ permutation matrix corresponding to π ; then $P_\pi w = \pi \cdot w$ for each $w \in \mathbb{R}^{\mathcal{M}}$. For any $x \in \mathbb{R}$, let $A(x)$ and $A^Q(x)$ be the $\mathcal{M} \times \mathcal{M}$ matrices defined by

$$(A(x))_{M,N} := x^{c(M,N)} \quad \text{and} \quad A^Q(x) := \sum_{s \in Q} A(x) P_s,
 \tag{16}$$

for $M, N \in \mathcal{M}$. So, by the weak reflection positivity of f , (15) implies that $A^Q(f(\bigcirc))$ is positive semidefinite. Note that each P_π commutes with $A(x)$, as for all $M, N \in \mathcal{M}$ one has $c(\pi \cdot M, \pi \cdot N) = c(M, N)$, implying $A(x) = P_\pi^T A(x) P_\pi = P_\pi^{-1} A(x) P_\pi$.

Hanlon and Wales [3] showed that the eigenvalues and eigenvectors of $A(x)$ can be described as follows. Consider any partition $\lambda = (t_1, \dots, t_m)$ of $8k$, with all t_i even. Then $A(x)$ has an eigenvalue

$$\mu_\lambda(x) := \prod_{a=1}^m \prod_{b=1}^{\frac{1}{2}t_a} (x - a + 2b - 1).
 \tag{17}$$

To describe a corresponding eigenvector, make a Young tableau T associated to λ such that each row of T has the form

$$\boxed{i_1 \mid i_1 \mid i_2 \mid i_2 \mid \dots \mid i_t \mid i_t}
 \tag{18}$$

for some $i_1, \dots, i_t \in [4k]$, where $\bar{i} := 4k + i$ for each $i \in [4k]$. For $i = 1, \dots, t_1$, let K_i denote the set of numbers in column i of T and let C_i be the subgroup of S_{8k} that permutes the elements of K_i . Then $C := C_1 \cdots C_{t_1}$. Similarly, for $i = 1, \dots, m$, let R_i be the subgroup of S_{8k} that permutes the numbers in row i of T , and $R := R_1 \cdots R_m$.

Let F be the perfect matching on $[8k]$ with edges $\{i, \bar{i}\}$ for $i \in [4k]$. Then

$$v := \sum_{c \in C, r \in R} \text{sgn}(c) cr \cdot F \tag{19}$$

is an eigenvector of $A(x)$ belonging to $\mu_\lambda(x)$. Then for $u := \sum_{q \in Q} q \cdot v$ one has

$$A^Q(x)u = \sum_{q', q \in Q} A(x)P_{q'}P_q v = \sum_{q', q \in Q} P_{q'}P_q A(x)v = \mu_\lambda(x) \sum_{q', q \in Q} P_{q'}P_q v = |Q|\mu_\lambda(x)u. \tag{20}$$

So u is an eigenvector of $A^Q(x)$ belonging to $|Q|\mu_\lambda(x)$, provided that u is nonzero. For this it suffices that the coefficient u_F of u in F is nonzero. Note that

$$u_F = \sum_{q \in Q} (q \cdot v)_F = \sum_{q \in Q} \sum_{c \in C, r \in R} \text{sgn}(c)(qcr \cdot F)_F = \sum_{\substack{q \in Q, c \in C, r \in R \\ qcr \cdot F = F}} \text{sgn}(c). \tag{21}$$

So $u \neq 0$ if for any $q \in Q, c \in C$, and $r \in R$, if $qcr \cdot F = F$ then $\text{sgn}(c) = 1$; that is (as Q is a group), if for any $q \in Q, c \in C, r \in R$:

$$\text{if } q \cdot F = cr \cdot F, \text{ then } \text{sgn}(c) = 1. \tag{22}$$

II. We first apply part I to the case where $f(\circ) \geq 0$. Let $k := \lceil f(\circ) \rceil + 1$, and consider the partition $\lambda := (8, 8, \dots, 8)$ of $8k$. Then, by (17),

$$\mu_\lambda(x) = \prod_{i=0}^{k-1} (x - i)(x - i + 2)(x - i + 4)(x - i + 6). \tag{23}$$

We give a Young tableau associated to λ that will yield (22). This implies that $|Q|\mu_\lambda(x)$ is an eigenvalue of $A^Q(x)$. So $\mu_\lambda(f(\circ)) \geq 0$. Hence, as the polynomial $\mu_\lambda(x)$ has largest zero $k - 1$, with multiplicity 1, and as $k - 1 = \lceil f(\circ) \rceil$, we know $f(\circ) = k - 1$.

Consider the following Young tableau associated to λ :

$$T := \begin{array}{|c|c|c|c|c|c|c|} \hline 1 & \bar{1} & 2 & \bar{2} & 3 & \bar{3} & 4 & \bar{4} \\ \hline 5 & \bar{5} & 6 & \bar{6} & 7 & \bar{7} & 8 & \bar{8} \\ \hline \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \hline 4k-3 & \overline{4k-3} & 4k-2 & \overline{4k-2} & 4k-1 & \overline{4k-1} & 4k & \overline{4k} \\ \hline \end{array}. \tag{24}$$

To prove (22), choose $q \in Q, c \in C$, and $r \in R$ with $q \cdot F = cr \cdot F$. Let $c = c_1 \cdots c_8$ with $c_i \in C_i$ ($i = 1, \dots, 8$) and define $M := q \cdot F$. Since F has no edges between $X := K_1 \cup K_2 \cup K_5 \cup K_6$ (the set of odd numbers in T) and $Y := K_3 \cup K_4 \cup K_7 \cup K_8$ (the set of even numbers in T) and since $Q \cdot X = X$ and $Q \cdot Y = Y$, we know that M has no edges between X and Y . For any $N \in \mathcal{M}$ and $Z \subseteq [8k]$, let N_Z be the set of edges of N contained in Z .

Let $z \in S_{8k}$ be defined by $z(i) := i + 1$ if 4 does not divide i and $z(i) := i - 3$ if 4 divides i . So $z^4 = \text{id}$, $z(X) = Y$, and $z \cdot F = F$. Moreover, $zq = qz$ (since $zb = bz$ and $zd = dz$ for all $b \in B$ and $d \in D$). So $z \cdot M = M$. Hence $z \cdot M_X = M_Y$.

Let $N := r \cdot F$. So $M = c \cdot N$. As no edge of M connects X and Y , also no edge in N connects X and Y . Moreover, as $z \cdot M_X = M_Y$, for each two columns K_i and K_j in X , we have $|M_{K_i \cup K_j}| = |M_{K_{i+2} \cup K_{j+2}}|$, and hence $|N_{K_i \cup K_j}| = |N_{K_{i+2} \cup K_{j+2}}|$. Moreover, if an edge $e \in N$ connects K_i and K_j , then N has an edge in the same row as e connecting the other two columns in X ; similarly for Y .

This implies that there exists a permutation $c' \in C_1 C_2 C_5 C_6$ that permutes complete rows in X in such a way that $c' \cdot N_X$ is a shift of N_Y ; that is, $zc' \cdot N_X = N_Y$. As c' maintains rows in X , there exists $r' \in R$ with $c' \cdot N = r' \cdot F$; so $c(c')^{-1}r' \cdot F = cr \cdot F$. Moreover, $\text{sgn}(c') = 1$, and, setting $N' := r' \cdot F$ we have $z \cdot N'_X = z \cdot (r' \cdot F)_X = z \cdot (c' \cdot N)_X = zc' \cdot N_X = N_Y = N'_Y$. Therefore, by replacing r by r' and c by $c(c')^{-1}$ we can assume that $z \cdot N_X = N_Y$.

Next consider any two columns K_i and K_j in X . Let $X' := K_i \cup K_j$ and $Y' := K_{i+2} \cup K_{j+2}$. So $Y' = z(X')$ and $z \cdot N_{X'} = N_{Y'}$. Then $e \mapsto z^{-1}c^{-1}zc(e)$ is a permutation σ of the edges e in $N_{X'}$, since $e \in N_{X'} \Rightarrow c(e) \in M_{X'} \Rightarrow zc(e) \in M_{Y'} \Rightarrow c^{-1}zc(e) \in N_{Y'} \Rightarrow z^{-1}c^{-1}zc(e) \in z^{-1} \cdot N_{Y'} = N_{X'}$. As σ permutes edges in X' , there exists a permutation $c' \in C_i C_j$ such that $c'(e) = z^{-1}c^{-1}zc(e)$ for all $e \in N_{X'}$ and such that c' only permutes elements covered by $N_{X'}$. Then $\text{sgn}(c') = 1$. By replacing c by $c(c')^{-1}$ we attain that $e = z^{-1}c^{-1}zc(e)$ for all edges $e \in N_{X'}$. So $cz(e) = zc(e)$ for all $e \in N_{X'}$.

Doing this for all K_i and K_j in X , we finally achieve that $cz(e) = zc(e)$ for all $e \in N_X$. As N_X is a perfect matching on X , this implies $cz(i) = zc(i)$ for all $i \in X$. Equivalently, $c_3 c_4 c_7 c_8 z(i) = z c_1 c_2 c_5 c_6(i)$ for all $i \in X$. Hence $\text{sgn}(c_3 c_4 c_7 c_8) = \text{sgn}(c_1 c_2 c_5 c_6)$, implying $\text{sgn}(c) = 1$.

III. Next we apply part I of this proof to the case where $f(\bigcirc) \leq 0$. Choose $k \in \mathbb{Z}_+$, and consider the partition $\lambda := (8k)$ of $8k$ and the following Young tableau

$$T := \begin{array}{|c|c|c|c|c|c|c|c|} \hline 1 & \bar{1} & 2 & \bar{2} & \dots & 4k-1 & \overline{4k-1} & 4k & \overline{4k} \\ \hline \end{array} \quad (25)$$

Then by (17),

$$\mu_\lambda(x) = \prod_{b=1}^{4k} (x - 2 + 2b). \quad (26)$$

Moreover, (22) trivially holds, as C only consists of the identity. The zeros of μ_λ are $-8k + 2, -8k + 4, -8k + 6, \dots, -2, 0$, all with multiplicity 1, so that $\mu_\lambda(f(\bigcirc)) \geq 0$ implies that $f(\bigcirc)$ does not belong to any interval $(-4t - 2, -4t)$ for any $t \in \mathbb{Z}_+$ with $t < 2k$. As k can be chosen arbitrarily large, we know that $f(\bigcirc) \notin (-4t - 2, -4t)$ for all $t \in \mathbb{Z}_+$.

To exclude the intervals $(-4t - 4, -4t - 2)$, consider the partition $\lambda := (8k - 2, 2)$ of $8k$ and the Young tableau

$$T := \begin{array}{|c|c|c|c|c|c|c|c|} \hline 1 & \bar{1} & 3 & \bar{3} & 4 & \bar{4} & \dots & 4k-1 & \overline{4k-1} & 4k & \overline{4k} \\ \hline 2 & \bar{2} & & & & & & & & & \\ \hline \end{array} \quad (27)$$

In this case, by (17),

$$\mu_\lambda(x) = (x - 1) \prod_{b=1}^{4k-1} (x - 2 + 2b). \quad (28)$$

To show (22), let $c = c_1 c_2$ with $c_1 \in C_1, c_2 \in C_2$. Observe that $M := q \cdot F$ contains no edges connecting an odd number with an even number (as F does not, and as Q maintains the sets of odd and even numbers).

If $\{2, \bar{2}\}$ belongs to M , then either c_1 and c_2 both are the identity permutation, or c_1 and c_2 both are transpositions. In either case, $\text{sgn}(c) = 1$ follows.

If $\{2, \bar{2}\}$ does not belong to M , then 2 and $\bar{2}$ are matched in M to even numbers in the first row of T . In this case, both c_1 and c_2 are transpositions, and again $\text{sgn}(c) = 1$ follows. This proves (22).

Now the zeros of μ_λ are $-8k + 4, -8k + 6, \dots, -2, 0, 1$, all with multiplicity 1, so that, like above, $f(\bigcirc) \notin (-4t - 4, -4t - 2)$ for all $t \in \mathbb{Z}_+$. □

4 Proof of the theorem

To see necessity in the theorem, let R be an R-matrix, say $R \in \mathcal{R}_n$. Then f_R is trivially multiplicative. Positive semidefiniteness of $M_{f_R, k}$ follows from

$$f_R(G \overset{k}{\vee} H) = p_n(G \overset{k}{\vee} H)(R) = \langle d^k p_n(G)(R), d^k p_n(H)(R) \rangle, \tag{29}$$

using (12).

To prove sufficiency, let f satisfy the conditions of the theorem. As $f(\bigcirc) \geq 0$ by assumption, the lemma implies that $n := f(\bigcirc)$ is a nonnegative integer. Then

$$\text{there exists an algebra homomorphism } F : p_n(\mathbb{R}\mathcal{G}) \rightarrow \mathbb{R} \text{ such that } f = F \circ p_n. \tag{30}$$

Otherwise, as f and p_n are algebra homomorphisms, there exists a quantum virtual link diagram γ with $p_n(\gamma) = 0$ and $f(\gamma) \neq 0$. We can assume that $p_n(\gamma)$ is homogeneous, that is, all virtual link diagrams in γ have the same number of crossings, k say. So $\gamma \overset{k}{\vee} \gamma$ has no crossings, that is, it is a polynomial in \bigcirc . As moreover $f(\bigcirc) = n = p_n(\bigcirc)$, we have $f(\gamma \overset{k}{\vee} \gamma) = p_n(\gamma \overset{k}{\vee} \gamma) = 0$, the latter equality because of (12). Similarly to Lemma 1 of [9], γ belongs to the ideal in $\mathbb{R}\mathcal{G}$ generated by $\gamma \overset{k}{\vee} \beta^i$ ($i = 0, \dots, k$), where β is the virtual link diagram

$$\beta := \left[\text{diagram of two linked loops} \right]. \tag{31}$$

(Note that $G \overset{1}{\vee} \beta = 2|V(G)|G$ for each virtual link diagram G .) As $f(\gamma \overset{k}{\vee} \gamma) = 0$ implies that $f(\gamma \overset{k}{\vee} \beta^i) = 0$ for each i (by the weak reflection positivity of f), we know $f(\gamma) = 0$, proving (30).

Now, by (13), $p_n(\mathbb{R}\mathcal{G}) = \mathcal{O}(\mathcal{R}_n)^{\mathcal{O}(n)}$. Basic invariant theory then gives the existence of an R in the complex extension of \mathcal{R}_n such that $F(q) = q(R)$ for each $q \in \mathcal{O}(\mathcal{R}_n)^{\mathcal{O}(n)}$ (cf. [9]). To prove that we can take R real, we apply the Procesi–Schwarz theorem [8].

For all $G, H \in \mathcal{G}$, using (12):

$$F(\langle dp_n(G), dp_n(H) \rangle) = F(p_n(G \overset{1}{\vee} H)) = f(G \overset{1}{\vee} H) = (M_{f,1})_{G,H}. \tag{32}$$

Since $M_{f,1}$ is positive semidefinite, (32) implies $F(\langle dq, dq \rangle) \geq 0$ for each $q \in p_n(\mathbb{R}\mathcal{G}) = \mathcal{O}(\mathcal{R}_n)^{\mathcal{O}(n)}$. Then by [8] there exists a (real) $R \in \mathcal{R}_n$ such that $F(q) = q(R)$ for each $q \in \mathcal{O}(\mathcal{R}_n)^{\mathcal{O}(n)} = p_n(\mathbb{R}\mathcal{G})$. Then $f = f_R$, as $f(G) = F(p_n(G)) = p_n(G)(R) = f_R(G)$ for each $G \in \mathcal{G}$.

One may finally check that substituting $f := f_R$ in (9), condition (9) (i) is equivalent to

$$\sum_{i,j} \left(\sum_a R_{iaaj} - \delta_{ij} \right)^2 = 0, \tag{33}$$

and hence to (6) (i); condition (9) (ii) is equivalent to

$$\sum_{i,j,k,l} \left(\sum_{a,b} R_{ijab} R_{alkb} - \delta_{ik} \delta_{jl} \right)^2 = 0, \tag{34}$$

and hence to (6) (ii); and condition (9) (iii) is equivalent to

$$\sum_{i,j,k,l,m,h} \left(\sum_{a,b,c} R_{iabh} R_{jkca} R_{bclm} - \sum_{a,b,c} R_{ijbc} R_{bkla} R_{camh} \right)^2 = 0, \tag{35}$$

and hence to (6) (iii). So R is an R-matrix, as required. □

References

1. Freedman, M.H., Lovász, L., Schrijver, A.: Reflection positivity, rank connectivity, and homomorphisms of graphs. *J. Am. Math. Soc.* **20**, 37–51 (2007)
2. Goodman, R., Wallach, N.R.: *Symmetry, Representations, and Invariants*. Springer, Dordrecht (2009)
3. Hanlon, P., Wales, D.: On the decomposition of Brauer’s centralizer algebras. *J. Algebra* **121**, 409–445 (1989)
4. de la Harpe, P., Jones, V.F.R.: Graph invariants related to statistical mechanical models: examples and problems. *J. Comb. Theory Ser. B* **57**, 207–227 (1993)
5. Kauffman, L.H.: Virtual knot theory. *Eur. J. Comb.* **20**, 663–690 (1999)
6. Kauffman, L.H.: Introduction to virtual knot theory. *J. Knot Theory Ramif.* **21**, 1240007 (2012)
7. Manturov, V.O., Ilyutko, D.P.: *Virtual Knots—The State of the Art*. World Scientific, River Edge (2013)
8. Procesi, C., Schwarz, G.: Inequalities defining orbit spaces. *Invent. Math.* **81**, 539–554 (1985)
9. Regts, G., Schrijver, A., Sevenster, B.: On partition functions for 3-graphs. *J. Combin. Theory Ser. B.* **121**, 421–431 (2016)
10. Sagan, B.E.: *The Symmetric Group: Representations, Combinatorial Algorithms, and Symmetric Functions*. Graduate Texts in Mathematics, vol. 203. Springer, New York (2001)
11. Schrijver, A.: On virtual link invariants (2012). [arXiv:1211.3572](https://arxiv.org/abs/1211.3572)
12. Szegedy, B.: Edge coloring models and reflection positivity. *J. Am. Math. Soc.* **20**, 969–988 (2007)