

# Survey of Problems, Questions, and Conjectures

We here collect unsolved problems, questions, and conjectures mentioned in this book. For terminology and background, we refer to the pages indicated.

**1** (page 41). Is  $NP \neq P$ ?

**2** (page 42). Is  $P = NP \cap \text{co-NP}$ ?

**3** (page 65). The *Hirsch conjecture*: A  $d$ -dimensional polytope with  $m$  facets has diameter at most  $m - d$ .

**4** (page 161). Is there an  $O(nm)$ -time algorithm for finding a maximum flow?

**5** (page 232). Berge [1982b] posed the following conjecture generalizing the Gallai-Milgram theorem. Let  $D = (V, A)$  be a digraph and let  $k \in \mathbb{Z}_+$ . Then for each path collection  $\mathcal{P}$  partitioning  $V$  and minimizing

$$(1) \quad \sum_{P \in \mathcal{P}} \min\{|VP|, k\},$$

there exist disjoint stable sets  $C_1, \dots, C_k$  in  $D$  such that each  $P \in \mathcal{P}$  intersects  $\min\{|VP|, k\}$  of them. This was proved by Saks [1986] for acyclic graphs.

**6** (page 403). The following open problem was mentioned by Fulkerson [1971b]: Let  $\mathcal{A}$  and  $\mathcal{B}$  be families of subsets of a set  $S$  and let  $w \in \mathbb{Z}_+^S$ . What is the maximum number  $k$  of common transversals  $T_1, \dots, T_k$  of  $\mathcal{A}$  and  $\mathcal{B}$  such that

$$(2) \quad \chi^{T_1} + \dots + \chi^{T_k} \leq w?$$

**7** (page 459). Can the weighted matching problem be formulated as a linear programming problem of size bounded by a polynomial in the size of the graph, by extending the set of variables? That is, is the matching polytope of a graph  $G = (V, E)$  equal to the projection of some polytope  $\{x \mid Ax \leq b\}$  with  $A$  and  $b$  having size bounded by a polynomial in  $|V| + |E|$ ?

**8** (pages 472,646). The *5-flow conjecture* of Tutte [1954a]:

$$(3) \quad (?) \text{ each bridgeless graph has a nowhere-zero 5-flow. } (?)$$

(A *nowhere-zero  $k$ -flow* is a flow over  $\mathbb{Z}_k$  in some orientation of the graph, taking value 0 nowhere.)

**9** (pages 472,498,645,1426). The *4-flow conjecture* of Tutte [1966]:

- (4) (?) each bridgeless graph without Petersen graph minor has a nowhere-zero 4-flow. (?)

This implies the four-colour theorem. For cubic graphs, (4) was proved by Robertson, Seymour, and Thomas [1997], Sanders, Seymour, and Thomas [2000], and Sanders and Thomas [2000].

Seymour [1981c] showed that the 4-flow conjecture is equivalent to the following more general conjecture, also due to Tutte [1966]:

- (5) (?) each bridgeless matroid without  $F_7^*$ ,  $M^*(K_5)$ , or  $M(\mathbf{P}_{10})$  minor has a nowhere-zero flow over  $\text{GF}(4)$ . (?)

Here  $\mathbf{P}_{10}$  denotes the Petersen graph.

**10** (page 472). The *3-flow conjecture* (W.T. Tutte, 1972 (cf. Bondy and Murty [1976], Unsolved problem 48)):

- (6) (?) each 4-edge-connected graph has a nowhere-zero 3-flow. (?)

**11** (page 473). The *weak 3-flow conjecture* of Jaeger [1988]:

- (7) (?) there exists a number  $k$  such that each  $k$ -edge-connected graph has a nowhere-zero 3-flow. (?)

**12** (page 473). The following *circular flow conjecture* of Jaeger [1984] generalizes both the 3-flow and the 5-flow conjecture:

- (8) (?) for each  $k \geq 1$ , any  $4k$ -connected graph has an orientation such that in each vertex, the indegree and the outdegree differ by an integer multiple of  $2k + 1$ . (?)

**13** (pages 475,645). The *generalized Fulkerson conjecture* of Seymour [1979a]:

- (9) (?)  $\lceil \chi'^*(G) \rceil = \lceil \frac{1}{2} \chi'(G_2) \rceil$  (?)

for each graph  $G$ . (Here  $\chi'^*(G)$  denotes the fractional edge-colouring number of  $G$ , and  $G_2$  the graph obtained from  $G$  by replacing each edge by two parallel edges.) This is equivalent to the conjecture that

- (10) (?) for each  $k$ -graph  $G$  there exists a family of  $2k$  perfect matchings, covering each edge precisely twice. (?)

(A  *$k$ -graph* is a  $k$ -regular graph  $G = (V, E)$  with  $|\delta(U)| \geq k$  for each odd-size subset  $U$  of  $V$ .)

**14** (pages 476,645). Fulkerson [1971a] asked if in each bridgeless cubic graph there exist 6 perfect matchings, covering each edge precisely twice (the *Fulkerson conjecture*). It is a special case of Seymour's generalized Fulkerson conjecture.

**15** (page 476). Berge [1979a] conjectures that the edges of any bridgeless cubic graph can be covered by 5 perfect matchings. (This would follow from the Fulkerson conjecture.)

**16** (page 476). Gol'dberg [1973] and Seymour [1979a] conjecture that for each (not necessarily simple) graph  $G$  one has

$$(11) \quad (?) \chi'(G) \leq \max\{\Delta(G) + 1, \lceil \chi^*(G) \rceil\}. (?)$$

An equivalent conjecture was stated by Andersen [1977].

**17** (page 476). Seymour [1981c] conjectures the following generalization of the four-colour theorem:

$$(12) \quad (?) \text{ each planar } k\text{-graph is } k\text{-edge-colourable. } (?)$$

For  $k = 3$ , this is equivalent to the four-colour theorem. For  $k = 4$  and  $k = 5$ , it was derived from the case  $k = 3$  by Guenin [2002b].

**18** (pages 476,644). Lovász [1987] conjectures more generally:

$$(13) \quad (?) \text{ each } k\text{-graph without Petersen graph minor is } k\text{-edge-colourable. } (?)$$

This is equivalent to stating that the incidence vectors of perfect matchings in a graph without Petersen graph minor, form a Hilbert base.

**19** (page 481). The following question was asked by Vizing [1968]: Is there a simple planar graph of maximum degree 6 and with edge-colouring number 7?

**20** (page 481). Vizing [1965a] asked if a minimum edge-colouring of a graph can be obtained from an arbitrary edge-colouring by iteratively swapping colours on a colour-alternating path or circuit and deleting empty colours.

**21** (page 482). Vizing [1976] conjectures that the list-edge-colouring number of any graph is equal to its edge-colouring number.

(The *list-edge-colouring number*  $\chi^l(G)$  of a graph  $G = (V, E)$  is the minimum number  $k$  such that for each choice of sets  $L_e$  for  $e \in E$  with  $|L_e| = k$ , one can select  $l_e \in L_e$  for  $e \in E$  such that for any two incident edges  $e, f$  one has  $l_e \neq l_f$ .)

**22** (page 482). Behzad [1965] and Vizing [1968] conjecture that the total colouring number of a simple graph  $G$  is at most  $\Delta(G) + 2$ . (The *total colouring*

*number* of a graph  $G = (V, E)$  is a colouring of  $V \cup E$  such that each colour consists of a stable set and a matching, vertex-disjoint.)

**23** (page 482). More generally, Vizing [1968] conjectures that the total colouring number of a graph  $G$  is at most  $\Delta(G) + \mu(G) + 1$ , where  $\mu(G)$  is the maximum edge multiplicity of  $G$ .

**24** (pages 497,645). Seymour [1979b] conjectures that each even integer vector in the circuit cone of a graph is a nonnegative integer combination of incidence vectors of circuits.

**25** (pages 497,645,1427). A special case of this is the *circuit double cover conjecture* (asked by Szekeres [1973] and conjectured by Seymour [1979b]): each bridgeless graph has circuits such that each edge is covered by precisely two of them.

Jamshy and Tarsi [1989] proved that the circuit double cover conjecture is equivalent to a generalization to matroids:

(14) (?) each bridgeless binary matroid without  $F_7^*$  minor has a circuit double cover. (?)

**26** (page 509). Is the system of  $T$ -join constraints totally dual quarter-integral?

**27** (page 517). L. Lovász asked for the complexity of the following problem: given a graph  $G = (V, E)$ , vertices  $s, t \in V$ , and a length function  $l : E \rightarrow \mathbb{Q}$  such that each circuit has nonnegative length, find a shortest odd  $s - t$  path.

**28** (page 545). What is the complexity of deciding if a given graph has a 2-factor without circuits of length at most 4?

**29** (page 545). What is the complexity of finding a maximum-weight 2-factor without circuits of length at most 3?

**30** (page 646). Tarsi [1986] mentioned the following strengthening of the circuit double cover conjecture:

(15) (?) in each bridgeless graph there exists a family of at most 5 cycles covering each edge precisely twice. (?)

**31** (page 657). Is the dual of any algebraic matroid again algebraic?

**32** (page 892). A special case of a question asked by A. Frank (cf. Schrijver [1979b], Frank [1995]) amounts to the following:

(16) (?) Let  $G = (V, E)$  be an undirected graph and let  $s \in V$ . Suppose that for each vertex  $t \neq s$ , there exist  $k$  internally vertex-disjoint  $s - t$  paths. Then  $G$  has  $k$  spanning trees such that for each vertex

$t \neq s$ , the  $s - t$  paths in these trees are internally vertex-disjoint.  
(?)

(The spanning trees need not be edge-disjoint — otherwise  $G = K_3$  would form a counterexample.) For  $k = 2$ , (16) was proved by Itai and Rodeh [1984, 1988], and for  $k = 3$  by Cheriyan and Maheshwari [1988] and Zehavi and Itai [1989].

**33** (page 962). Can a maximum number of disjoint directed cut covers in a directed graph be found in polynomial time?

**34** (page 962). Woodall [1978a,1978b] conjectures (*Woodall's conjecture*):

(17) (?) In a digraph, the minimum size of a directed cut is equal to the maximum number of disjoint directed cut covers. (?)

**35** (page 985). Let  $G = (V, E)$  be a complete undirected graph, and consider the system

$$(18) \quad \begin{aligned} 0 \leq x_e \leq 1 & \text{ for each edge } e, \\ x(\delta(v)) = 2 & \text{ for each vertex } v, \\ x(\delta(U)) \geq 2 & \text{ for each } U \subseteq V \text{ with } \emptyset \neq U \neq V. \end{aligned}$$

Let  $l : E \rightarrow \mathbb{R}_+$  be a length function. Is the minimum length of a Hamiltonian circuit at most  $\frac{4}{3}$  times the minimum value of  $l^T x$  over (18)?

**36** (page 990). Padberg and Grötschel [1985] conjecture that the diameter of the symmetric traveling salesman polytope of a complete graph is at most 2.

**37** (page 1076). Frank [1994a] conjectures:

(19) (?) Let  $D = (V, A)$  be a simple acyclic directed graph. Then the minimum size of a  $k$ -vertex-connector for  $D$  is equal to the maximum of  $\sum_{v \in V} \max\{0, k - \deg^{\text{in}}(v)\}$  and  $\sum_{v \in V} \max\{0, k - \deg^{\text{out}}(v)\}$ . (?)

(A  $k$ -vertex-connector for  $D$  is a set of (new) arcs whose addition to  $D$  makes it  $k$ -vertex-connected.)

**38** (page 1087). *Hadwiger's conjecture* (Hadwiger [1943]): If  $\chi(G) \geq k$ , then  $G$  contains  $K_k$  as a minor.

Hadwiger's conjecture is trivial for  $k = 1, 2, 3$ , was shown by Hadwiger [1943] for  $k = 4$  (also by Dirac [1952]), is equivalent to the four-colour theorem for  $k = 5$  (by a theorem of Wagner [1937a]), and was derived from the four-colour theorem for  $k = 6$  by Robertson, Seymour, and Thomas [1993]. For  $k \geq 7$ , the conjecture is unsettled.

**39** (page 1099). Chvátal [1973a] asked if for each fixed  $t$ , the stable set problem for graphs for which the stable set polytope arises from  $P(G)$  by at most

$t$  rounds of cutting planes, is polynomial-time solvable. Here  $P(G)$  is the polytope determined by the nonnegativity and clique inequalities.

**40** (page 1099). Chvátal [1975b] conjectures that there is no polynomial  $p(n)$  such that for each graph  $G$  with  $n$  vertices we can obtain the inequality  $x(V) \leq \alpha(G)$  from the system defining  $Q(G)$  by adding at most  $p(n)$  cutting planes. Here  $Q(G)$  is the polytope determined by the nonnegativity and edge inequalities. (This conjecture would be implied by  $\text{NP} \neq \text{co-NP}$ .)

**41** (page 1105). Gyárfás [1987] conjectures that there exists a function  $g : \mathbb{Z}_+ \rightarrow \mathbb{Z}_+$  such that  $\chi(G) \leq g(\omega(G))$  for each graph  $G$  without odd holes.

**42** (page 1107). Can perfection of a graph be tested in polynomial time?

**43** (page 1131). Berge [1982a] conjectures the following. A directed graph  $D = (V, A)$  is called  $\alpha$ -diperfect if for every induced subgraph  $D' = (V', A')$  and each maximum-size stable set  $S$  in  $D'$  there is a partition of  $V'$  into directed paths each intersecting  $S$  in exactly one vertex. Then for each directed graph  $D$ :

(20)            (?)  $D$  is  $\alpha$ -diperfect if and only if  $D$  has no induced subgraph  $C$  whose underlying undirected graph is a chordless odd circuit of length  $\geq 5$ , say with vertices  $v_1, \dots, v_{2k+1}$  (in order) such that each of  $v_1, v_2, v_3, v_4, v_6, v_8, \dots, v_{2k}$  is a source or a sink. (?)

**44** (page 1170). Is  $\vartheta(C_n) = \Theta(C_n)$  for each odd  $n$ ?

**45** (page 1170). Can Haemers' bound  $\eta(G)$  on the Shannon capacity of a graph  $G$  be computed in polynomial time?

**46** (page 1187). Is every  $t$ -perfect graph strongly  $t$ -perfect?

Here a graph is  $t$ -perfect if its stable set polytope is determined by the nonnegativity, edge, and odd circuit constraints. It is *strongly  $t$ -perfect* if this system is totally dual integral.

**47** (page 1195).  $T$ -perfection is closed under taking induced subgraphs and under contracting all edges in  $\delta(v)$  where  $v$  is a vertex not contained in a triangle. What are the minimally non- $t$ -perfect graphs under this operation?

**48** (page 1242). For any  $k$ , let  $f(k)$  be the smallest number such that in any  $f(k)$ -connected undirected graph, for any choice of distinct vertices  $s_1, t_1, \dots, s_k, t_k$  there exist vertex-disjoint  $s_1 - t_1, \dots, s_k - t_k$  paths. Thomassen [1980] conjectures that  $f(k) = 2k + 2$  for  $k \geq 2$ .

**49** (page 1242). For any  $k$ , let  $g(k)$  be the smallest number such that in any  $g(k)$ -edge-connected undirected graph, for any choice of vertices  $s_1, t_1, \dots, s_k, t_k$  there exist edge-disjoint  $s_1 - t_1, \dots, s_k - t_k$  paths. Thomassen [1980] conjectures that  $g(k) = k$  if  $k$  is odd and  $g(k) = k + 1$  if  $k$  is even.

- 50** (page 1243). What is the complexity of the  $k$  arc-disjoint paths problem in directed planar graphs, for any fixed  $k \geq 2$ ? This is even unknown for  $k = 2$ , also if we restrict ourselves to two opposite nets.
- 51** (page 1274). Karzanov [1991] conjectures that if the nets in a multifold problem form two disjoint triangles and if the capacities and demands are integer and satisfy the Euler condition, then the existence of a fractional multifold implies the existence of a half-integer multifold.
- 52** (page 1274). The previous conjecture implies that for each graph  $H = (T, R)$  without three disjoint edges, there is an integer  $k$  such that for each graph  $G = (V, E)$  with  $V \supseteq T$  and any  $c : E \rightarrow \mathbb{Z}_+$  and  $d : R \rightarrow \mathbb{Z}_+$ , if there is a feasible multifold, then there exists a  $\frac{1}{k}$ -integer multifold.
- 53** (page 1276). Okamura [1998] conjectures the following. Let  $G = (V, E)$  be an  $l$ -edge-connected graph (for some  $l$ ). Let  $H = (T, R)$  be a ‘demand’ graph, with  $T \subseteq V$ , such that  $d_R(U) \leq l$  for each  $U \subseteq V$ . Then the edge-disjoint paths problem has a half-integer solution.
- 54** (page 1293). Is each Mader matroid a gammoid?
- 55** (page 1294). Is each Mader matroid linear?
- 56** (page 1299). Is the undirected edge-disjoint paths problem for planar graphs polynomial-time solvable if all terminals are on the outer boundary? Is it NP-complete?
- 57** (page 1310). Is the integer multifold problem polynomial-time solvable if the graph and the nets form a planar graph such that the nets are spanned by a fixed number of faces?
- 58** (page 1310). Pfeiffer [1990] raised the question if the edge-disjoint paths problem has a half-integer solution if the graph  $G + H$  (the union of the supply graph and the demand graph) is embeddable in the torus and there exists a quarter-integer solution.
- 59** (page 1320). Let  $G = (V, E)$  be a planar bipartite graph and let  $q$  be a vertex on the outer boundary. Do there exist disjoint cuts  $C_1, \dots, C_p$  such that any pair  $s, t$  of vertices with  $s$  and  $t$  on the outer boundary, or with  $s = q$ , is separated by  $\text{dist}_G(s, t)$  cuts?
- 60** (page 1345). Fu and Goddyn [1999] asked: Is the class of graphs for which the incidence vectors of cuts form a Hilbert base, closed under taking minors?
- 61** (page 1382). Füredi, Kahn, and Seymour [1993] conjecture that for each hypergraph  $H = (V, \mathcal{E})$  and each  $w : \mathcal{E} \rightarrow \mathbb{R}_+$ , there exists a matching  $\mathcal{M} \subseteq \mathcal{E}$  such that

$$(21) \quad \sum_{F \in \mathcal{M}} \left( |F| - 1 + \frac{1}{|F|} \right) w(F) \geq \nu_w^*(H),$$

where  $\nu_w^*(H)$  is the maximum weight  $w^\top y$  of a fractional matching  $y : \mathcal{E} \rightarrow \mathbb{R}_+$ .

**62** (pages 1387,1408). Seymour [1981a] conjectures:

$$(22) \quad (?) \text{ a binary hypergraph is ideal if and only if it has no } \mathcal{O}(K_5), \\ b(\mathcal{O}(K_5)), \text{ or } F_7 \text{ minor. } (?)$$

**63** (page 1392). Seymour [1990b] asked the following. Suppose that  $H = (V, \mathcal{E})$  is a hypergraph without  $J_n$  minor ( $n \geq 3$ ). Let  $l, w : V \rightarrow \mathbb{Z}_+$  be such that

$$(23) \quad \tau(H^w) \cdot \tau(b(H)^l) > l^\top w.$$

Is there a minor  $H'$  of  $H$  and  $l', w' : V H' \rightarrow \{0, 1\}$  such that

$$(24) \quad \tau((H')^{w'}) \cdot \tau(b(H')^{l'}) > l'^\top w'$$

and such that  $\tau((H')^{w'}) \leq \tau(H^w)$  and  $\tau(b(H')^{l'}) \leq \tau(b(H)^l)$ ?

Here, for each  $n \geq 3$ :  $J_n :=$  the hypergraph with vertex set  $\{1, \dots, n\}$  and edges  $\{2, \dots, n\}, \{1, 2\}, \dots, \{1, n\}$ .

**64** (page 1392). Seymour [1990b] also asked the following. Let  $H = (V, \mathcal{E})$  be a nonideal hypergraph. Is the minimum of  $\tau(H')$  over all parallelizations and minors  $H'$  of  $H$  with  $\tau^*(H') < \tau(H')$  attained by a minor of  $H$ ?

**65** (page 1395). Cornuéjols and Novick [1994] conjecture that there are only finitely many minimally nonideal hypergraphs  $H$  with  $r_{\min}(H) > 2$  and  $\tau(H) > 2$ .

**66** (page 1396). Ding [1993] asked whether there exists a number  $t$  such that each minimally nonideal hypergraph  $H$  satisfies  $r_{\min}(H) \leq t$  or  $\tau(H) \leq t$ .

(The above conjecture of Cornuéjols and Novick [1994] implies a positive answer to this question.)

**67** (page 1396). Ding [1993] conjectures that for each fixed  $k \geq 2$ , each minor-minimal hypergraph  $H$  with  $\tau_k(H) < k \cdot \tau(H)$ , contains some  $J_n$  minor ( $n \geq 3$ ) or satisfies the regularity conditions of Lehman's theorems (Theorem 78.4 and 78.5).

**68** (page 1401). Conforti and Cornuéjols [1993] conjecture:

$$(25) \quad (?) \text{ a hypergraph is Mengerian if and only if it is packing. } (?)$$

**69** (page 1401). Cornuéjols, Guenin, and Margot [1998,2000] conjecture:

$$(26) \quad (?) \text{ each minimally nonideal hypergraph } H \text{ with } r_{\min}(H)\tau(H) = \\ |VH| + 1 \text{ is minimally nonpacking. } (?)$$

**70** (page 1401). Cornuéjols, Guenin, and Margot [1998,2000] conjecture that  $\tau(H) = 2$  for each ideal minimally nonpacking hypergraph  $H$ .

**71** (page 1404). Seymour [1981a] conjectures that  $T_{30}$  is the unique minor-minimal binary ideal hypergraph  $H$  with the property  $\nu_2(H) < 2\tau(H)$ .

Here the hypergraph  $T_{30}$  arises as follows. Replace each edge of the Petersen graph by a path of length 2, making the graph  $G$ . Let  $T := VG \setminus \{v\}$ , where  $v$  is an arbitrary vertex of  $v$  of degree 3. Let  $\mathcal{E}$  be the collection of  $T$ -joins. Then  $T_{30} := (EG, \mathcal{E})$ .

**72** (page 1405). P.D. Seymour (personal communication 1975) conjectures that for each ideal hypergraph  $H$  there exists an integer  $k$  such that  $\nu_k(H) = k \cdot \tau(H)$  and such that  $k = 2^i$  for some  $i$ . He also asks if  $k = 4$  would do in all cases.

**73** (page 1405). Seymour [1979a] conjectures that for each ideal hypergraph  $H$ , the g.c.d. of those  $k$  with  $\nu_k(H) = k \cdot \tau(H)$  is equal to 1 or 2.

**74** (page 1409). Is the following true for binary hypergraphs  $H$ :

$$(27) \quad \begin{aligned} (?) \nu(H^w) = \tau(H^w) \text{ for each } w : V \rightarrow \mathbb{Z}_+ \text{ with } w(B) \text{ even for all} \\ B \in b(H) \iff \frac{1}{2}\nu_2(H^w) = \tau(H^w) \text{ for each } w : V \rightarrow \mathbb{Z}_+ \iff \\ H \text{ has no } \mathcal{O}(K_5), b(\mathcal{O}(K_5)), F_7, \text{ or } T_{15} \text{ minor. } (?) \end{aligned}$$

Here  $T_{15}$  is the hypergraph of  $VP_{10}$ -joins in the Petersen graph  $\mathbf{P}_{10}$ .

**75** (page 1421). Seymour [1981a] conjectures that for any binary matroid  $M$ :

$$(28) \quad \begin{aligned} (?) M \text{ is 1-cycling} \iff M \text{ is 1-flowing} \iff M \text{ has no AG}(3,2), \\ T_{11}, \text{ or } T_{11}^* \text{ minor. } (?) \end{aligned}$$

Here  $T_{11}$  is the binary matroid represented by the 11 vectors in  $\{0,1\}^5$  with precisely 3 or 5 ones. Moreover,  $\text{AG}(3,2)$  is the matroid with 8 elements obtained from the 3-dimensional affine geometry over  $\text{GF}(2)$ .